


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1967

[Full Table of Content](#)



---

### **67/S/SAX/B/1/C/44 Formalin N and Lime - Barley**

#### **Rothamsted Research**

Rothamsted Research (1968) *67/S/SAX/B/1/C/44 Formalin N and Lime - Barley* ; Yields Of The Field Experiments 1967, pp 265 - 267 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/C/44.1

BARLEY

(SAX/B/1)

Formalin, nitrogen and lime for barley, Saxmundham, Grove Plot, 1967, the first year.

Design: A single replicate of 2 x 2 x 2 x 2 x 2 in 2 blocks of 16 plots, with 2 additional plots per block.

Area of each plot: 0.0022. Area harvested: 0.0011.

Treatments: All combinations of:-

1. Formalin: None (O), formalin at 532 gals in 2900 gals (F).
2. Nitrogen: 0.6 (N1), 1.2 (N2) cwt N as calcium nitrate.
3. Time of application of nitrogen: Early (E), late (L).
4. Lime: None (O), 7.5 tons of ground chalk (C).
5. Variety: Deba Abed (A), Maris Badger (B).

Additional plots: One of Deba Abed (A), one of Maris Badger (B), each with no formalin, nitrogen or lime.

Basal applications: 0.5 cwt P<sub>2</sub>O<sub>5</sub>, 0.5 cwt K<sub>2</sub>O as (0:20:20).

Weedkiller: 2,4-DP/MCPA (Cornox RK Extra at 6 pints in 50 gals).

Cultivations, etc.: Ploughed: Nov 29 - Dec 16, 1966.

Formalin applied: Feb 7, 1967. Basal PK and first dressing of ground chalk (5 tons) applied: Feb 8. Calcium nitrate (E treatment) and second dressing of ground chalk (2.5 tons) applied, seed drilled at 160 lb: Mar 21. Calcium nitrate (L treatment) and weedkiller applied: May 11. Harvested by hand: Aug 15.

NOTE: Soil samples were taken for N determination before sowing.

Standard error per plot.

Grain: 2.71 or 6.9% (5 d.f.)

67/C/44.2

SUMMARY OF RESULTS

GRAIN

	N1	N2	E	L	O	C	A	B	Mean
	(±0.96)		(±0.96)		(±0.96)		(±0.96)		(±0.68)
O	36.0	40.9	35.1	41.8	38.8	38.1	39.7	37.3	38.5
F	39.8	40.5	37.5	42.8	39.4	40.9	41.9	38.5	40.2
			(±0.96)		(±0.96)		(±0.96)		(±0.68)
		N1	31.6	44.2	37.4	38.5	38.1	37.7	37.9
		N2	41.0	40.4	40.8	40.6	43.4	38.0	40.7
					(±0.96)		(±0.96)		(±0.68)
				E	35.8	36.9	37.0	35.7	36.3
				L	42.4	42.2	44.6	40.0	42.3
							(±0.96)		(±0.68)
						O	40.6	37.6	39.1
						C	40.9	38.1	39.5
Mean (±0.68)							40.8	37.9	39.3

Additional plots

A	B
16.1	27.7

General mean: 37.4  
 Mean D.M. %: 79.1

67/C/4.3

		STRAW							Mean	
		N1	N2	E	L	O	C	A	B	
O		35.9	44.3	35.7	44.5	40.6	39.6	36.9	43.4	40.1
	F	40.4	45.0	38.8	46.6	41.3	44.1	40.4	44.9	42.7
			N1	31.4	44.9	37.6	38.7	34.5	41.8	38.1
			N2	43.0	46.3	44.2	45.1	42.8	46.6	44.7
				E		37.3	37.1	33.7	40.7	37.2
				L		44.5	46.7	43.5	47.6	45.6
							O	37.7	44.2	40.9
							C	39.6	44.1	41.9
Mean								38.6	44.2	41.4

Additional plots

A	B
13.5	28.5

General mean: 39.1  
 Mean D.M. %: 74.9