


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/S/SAX/G/1/C/47 N and Cutting - Grass

Rothamsted Research

Rothamsted Research (1968) *67/S/SAX/G/1/C/47 N and Cutting - Grass* ; Yields Of The Field Experiments 1967, pp 273 - 274 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/C/47.1

GRASS

(SAX/G/1)

N and cutting, Saxmundham Grove Plot 1967, the first year.

Design: 4 randomised blocks of 9 plots.

Area of each plot: 0.0018. Area harvested: 0.0013.

Treatments:- (N as 'Nitro-Chalk').

	Total N (cwt)	No. of N dressings	No. of cuts
1	0.0	0	2
2	1.0	2	2
3	2.0	2	2
4	3.0	2	2
5	4.0	2	2
6	1.5	3	3
7	3.0	3	3
8	2.0	2	3
9	4.0	2	3

Basal applications: 1.0 cwt P205, 2.0 cwt K20 as (0:10:20).

Seeds mixture lb per acre

S352 Early Timothy	95 lb
S215 Meadow Fescue	95 lb

Cultivations, etc.: Ploughed: Nov 29 - Dec 16, 1966. Basal PK applied: Mar 8, 1967. First N treatments applied, seed sown: Mar 22. 2nd and 3rd N treatments applied: June 28 and July 18. Cut plots with 2 cuts: July 18 and Sept 26, plots with 3 cuts: June 28, Aug 15, Sept 26. Previous crops: Barley 1965, fallow 1966.

Standard error per plot. Dry matter:

Total of all cuts: 3.80 or 7.0% (24 d.f.)

67/c/47.2

SUMMARY OF RESULTS

DRY MATTER. TOTAL OF ALL CUTS

1	2	3	4	5	6	7	8	9	Mean
(±1.90)									
24.1	49.2	57.2	61.9	67.6	48.6	61.7	51.4	64.7	54.0

Mean D.M. %: 2 cut plots: 26.0
3 cut plots: 20.7