

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/RW801/DA/8 and 67/W/WW301/DA/8 Spring Wheat - Anhydrous Ammonia

Rothamsted Research

Rothamsted Research (1968) *67/R/RW801/DA/8 and 67/W/WW301/DA/8 Spring Wheat - Anhydrous Ammonia* ; Yields Of The Field Experiments 1967, pp 289 - 291 - DOI:

<https://doi.org/10.23637/ERADOC-1-157>

67/Da/8.1

SPRING WHEAT

(RW801 and WW301)

Anhydrous ammonia as a fertiliser - Rothamsted (R), Great Knott I and Woburn (W) Great Hill S.W. 1967.

Design: 4 randomised blocks of 8 plots.

Area of each plot: 0.0321. Area harvested: 0.0213.

Treatments: All combinations of:-

1. Forms of N: Broadcast 'Nitro-Chalk' (B), injected anhydrous ammonia (I).
2. Levels of N: 0.5 (N1), 1.0 (N2), 1.5 (N3) cwt N together with no nitrogen - without (NO), with (NOI), the injector running idle through the soil.

Basal applications:

Great Knott I (R): 40 cwt ground chalk, 250 lb (0:20:20) combine drilled. Weedkiller: Mecoprop/2,4-D (Methoxone Extra at 6 pints in 30 gals).

Great Hill S.W. (W): 40 cwt ground chalk, 280 lb (0:20:20) combine drilled. Weedkiller: Ioxynil/mecoprop (Actril C at 5 pints in 35 gals).

Cultivations, etc.:

Great Knott I (R): Ground chalk applied: Nov 11, 1966. Ploughed: Sept 28. Anhydrous ammonia injected: Mar 8, 1967. 'Nitro-Chalk' applied, seed drilled at 185 lb: Mar 14. Weedkiller applied: May 11. Combine harvested: Aug 24. Variety: Kloka. Previous crops: Potatoes 1965, winter wheat 1966.

Great Hill S.W. (W): Ground chalk applied: Nov 3, 1966. Deep-tine cultivated: Feb 3, 1967. Anhydrous ammonia injected: Mar 8. 'Nitro-Chalk' applied: Mar 13. Seed drilled at 170 lb: Mar 14. Weedkiller applied: May 3. Combine harvested: Aug 22. Variety: Kloka. Previous crops: Barley 1965, potatoes 1966.

NOTES: (1) Green crop samples were taken on 28th June, for yield and N content.

(2) Owing to a failure of the injector little ammonia was injected into plot WW326 - IN2. An estimated value was used in the analysis.

Standard errors per plot (estimated from plots receiving N). Grain:

Great Knott I (R): 2.28 or 6.9% (15 d.f.)

Great Hill S.W.(W): 4.51 or 13.6% (14 d.f.)*

* 1 estimated value.

67/Da/8.2

SUMMARY OF RESULTS

GREAT KNOTT I (R)

GRAIN

	N1	N2	N3	Mean
		(±1.14)		(±0.66)
B	28.5	35.4	36.3	33.4
I	26.8	32.5	38.0	32.4
Mean (±0.81)	27.7	34.0	37.1	32.9

	NO	NOI	Mean
	18.8	20.1	19.4
	(±1.14)		

General mean: 29.6

Mean D.M. %: 80.9

67/De/8.3

GREAT HILL S.W. (W)

GRAIN

	N1	N2	N3	Mean
		(±2.25)		(±1.30)
B	30.7	35.6	42.5	36.3
I	20.1	33.3	36.4	29.9
Mean (±1.59)	25.4	34.5	39.5	33.1

	NO	NOI	Mean
	15.8	16.6	16.2
	(±2.25)		

General mean: 28.9

Mean D.M. %: 84.3