


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1967

[Full Table of Content](#)



### 67/R/BG1/DA/11 Winter Wheat - Sowing Date and Bulb Fly

#### Rothamsted Research

Rothamsted Research (1968) *67/R/BG1/DA/11 Winter Wheat - Sowing Date and Bulb Fly* ; Yields Of The Field Experiments 1967, pp 299 - 300 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/Da/11.1

WINTER WHEAT

(BG 1)

Sowing dates and bulb fly - Stackyard 1967, the third year.

Design: 3 randomised blocks of 3 plots, split into 2 for covering to prevent egg-laying (unrandomised).

Area of each sub plot: 0.0096. Area harvested: 0.0064.

Treatments: All combinations of:-

Whole plots: 1. Sowing dates: Oct 28 (E), Nov 29 (M),  
Dec 21 (L). (Seed rate 180 lb.)

Sub plots: 2. Not covered (O), covered with polythene  
sheet to prevent egg-laying, late June to  
mid-September 1966 (C).

Basal applications: 280 lb (6:15:15) combine drilled, 0.84 cwt N  
as 'Nitro-Chalk' in spring. Weedkillers: Diquat (Reglone at  
4 pints in 40 gals), ioxynil/mecoprop (Actril C at 6 pints in  
20 gals). Seed dressed with organo-mercury fungicide only.

Cultivations, etc.: Rotary cultivated twice: Mar 30 and May 19,  
1966. Deep-tine cultivated: June 9. Rotary cultivated:  
July 1. Diquat applied: Oct 26. 'Nitro-Chalk' applied:  
Apr 18, 1967. Ioxynil/mecoprop applied: May 9. Combine  
harvested: Aug 30. Variety: Cappelle. Previous crops:  
Winter wheat 1965, fallow 1966.

- NOTES: (1) For the previous year's results see 'Results'  
65/Da/5 and 66/Da/3.
- (2) Samples were taken from late February until mid-May  
to estimate numbers of plants, shoots and larvae  
and damaged plants and shoots. Samples were taken  
just before harvest to estimate ear number and grain  
weight.
- (3) It was intended to sow four blocks, but on one the  
soil was too wet.

Standard errors per plot. Grain:  
Whole plot: 6.46 or 15.9% (4 d.f.)  
Sub plot: 5.57 or 13.8% (6 d.f.)

67/Da/11.2

SUMMARY OF RESULTS

GRAIN

	E	M	L	Mean
	(1) and (2)			
D	49.6	39.5	27.4	38.8
C	50.6	42.8	33.5	42.3
Mean ( $\pm 3.73$ )	50.1	41.1	30.4	40.6

- (1) ( $\pm 5.27$ ) For use in horizontal comparisons only  
 (2) ( $\pm 3.22$ ) For use in interaction comparisons only

Mean D.M. %: 84.1