


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1967

[Full Table of Content](#)



---

### 67/R/RBE/101/DC/1 Spring Beans - *Rhizobium* Strains

#### Rothamsted Research

Rothamsted Research (1968) *67/R/RBE/101/DC/1 Spring Beans - Rhizobium Strains* ; Yields Of The Field Experiments 1967, pp 309 - 310 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/Dc/1.1

SPRING BEANS

(RBe 101)

Rhizobium strains - Long Hoos III 1967.

Design: 4 randomised blocks of 4 plots.

Area of each plot: 0.0202. Area harvested: 0.0127.

Treatments: Seed uninoculated (R0), inoculated with Rhizobium strain 1027 (R1), strain CZS (R2), strain 1007 (R3).

Basal applications: 360 lb (0:14:28) placement drilled. Weedkiller: Simazine at 1 lb a.i. in 32 gals. Insecticide: Demeton-s-methyl on two occasions at 3 oz in 37 gals, and at 3 oz in 3 gals by fixed-wing aircraft.

Cultivations, etc.: Ploughed: Sept 22, 1966. Ground chalk applied at 50 cwt: Jan 7, 1967. Ploughed second time: Jan 13. Seed drilled at 200 lb: Mar 16. Simazine applied: Mar 18. Insecticide applied: June 14 and July 11. Combine harvested: Sept 1. Variety: Tarvin. Previous crops: Potatoes 1965, spring wheat 1966.

NOTE: Counts of Rhizobium leguminosarum in the soil were made in March before sowing and from the rhizosphere and soil of selected plots in May, using hairy vetch (*Vicia hirsuta*). Counts of plants were taken in August.

Standard error per plot.

Grain: 1.43 or 4.2% (9 d.f.)

67/Dc/1.2

SUMMARY OF RESULTS

GRAIN

R0	R1	R2	R3	Mean
34.2	34.9	33.9	34.7	34.4
(±0.71)				

Mean D.M. %: 79.2