

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1967

[Full Table of Content](#)



---

### 67/R/RP5/1/DD/4 Potatoes - Rhizoctonia

#### Rothamsted Research

Rothamsted Research (1968) *67/R/RP5/1/DD/4 Potatoes - Rhizoctonia* ; Yields Of The Field Experiments 1967, pp 319 - 320 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/Dd/4.1

POTATOES

(RP 5/1)

Effects of stem-canker (*Rhizoctonia solani*) - Pastures 1967.

Design: 6 randomised blocks of 2 plots, split into 4. 3 blocks with weedkiller, 3 without (not randomised).

Area of each sub plot: 0.0036. Area harvested: 0.0033.

Treatments: All combinations of:-

Columns of

3 blocks: 1. Weedkiller: None, with inter-row cultivation (O), sprayed with linuron at 0.75 lb and paraquat at 0.5 lb ion in 37 gals, no inter-row cultivation (W).

Plots: 2. Varieties: King Edward (E), Majestic (M).

Sub plots: 3. Levels of seed infection: Clean (A), moderately infected (B), severely infected (C), unselected stock (D).

Basal applications: 10 cwt (13:13:20). Fungicide: Mancozeb at 1.2 lb in 30 gals on four occasions.

Cultivations, etc.: Rotary cultivated: Sept 16, 1966. Ploughed: Nov 23 - Jan 5, 1967. Basal compound fertiliser applied: Mar 30. Rotary cultivated, potatoes planted: Apr 19. Weedkiller applied: May 17. 'O' blocks grubbed and mechanically weeded: May 19. 'O' blocks grubbed: June 14. Fungicide applied: June 30, July 21, Aug 3, Aug 24. Sprayed with undiluted BOV at 15 gals: Sept 20. Haulm destroyed mechanically: Sept 25. Lifted: Sept 28 and Oct 5. Previous crops: Spring wheat and barley 1965, fallow 1966.

Standard errors per plot. Total tubers:

Whole plots: 0.812 or 3.8% (4 d.f.)

Sub plots: 1.866 or 8.8% (24 d.f.)

67/Dd/4.2

SUMMARY OF RESULTS

	E	M	A	B	C	D	Mean
TOTAL TUBERS							
	( $\pm 0.469$ )*		( $\pm 0.762$ )*				
O	19.36	20.63	20.35	21.24	18.30	20.09	20.00
W	22.56	22.73	23.75	22.01	22.40	22.44	22.65
			(1) and (2)				( $\pm 0.332$ )
	E		21.48	21.78	19.90	20.70	20.96
	M		22.62	21.47	20.80	21.82	21.68
Mean ( $\pm 0.539$ )			22.05	21.62	20.35	21.26	21.32

\* For use in horizontal and interaction comparisons only

(1) ( $\pm 0.738$ ) For use in vertical and diagonal comparisons

(2) ( $\pm 0.762$ ) For use in horizontal and interaction comparisons

% WARE							
O	82.4	90.0	85.2	87.6	84.1	87.9	86.2
W	84.9	90.9	88.2	88.1	87.0	88.3	87.9
	E		84.3	84.9	80.5	84.9	83.6
	M		89.1	90.8	90.5	91.3	90.5
Mean			86.7	87.9	85.5	88.1	87.0