


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/RP10/1/DD/9 and 67/W/WP1/1/DD/9 Potatoes - Chitting and Scab

Rothamsted Research

Rothamsted Research (1968) *67/R/RP10/1/DD/9 and 67/W/WP1/1/DD/9 Potatoes - Chitting and Scab* ; Yields Of The Field Experiments 1967, pp 331 - 332 - DOI:

<https://doi.org/10.23637/ERADOC-1-157>

67/Dd/9.1

POTATOES

(RP 10/1 and WP 101)

Chitting and scab - Rothamsted (R) Pastures and Woburn (W) Great Hill Bottom I, 1967.

Design: 4 randomised blocks of 3 plots.

Area of each plot: 0.0150. Area harvested: 0.0007.

Treatments:

Non-chitted seed planted Apr 20 - 21 (OE), May 18 (OL), chitted seed planted Apr 20 - 21 (CE).

Basal applications: 10 cwt (13:13:20). Weedkiller: Linuron at 0.75 lb plus paraquat at 0.5 lb in 37 gals. Fungicides:- Pastures (R): Mancozeb at 1.2 lb in 30 gals. Great Hill (W): Fentin acetate at 4.2 oz plus maneb at 1.4 lb in 30 gals on three occasions, and mancozeb at 1.2 lb in 30 gals.

Cultivations, etc.:

Pastures (R): Rotary cultivated: Sept 16, 1966. Ground chalk applied at 50 cwt: Jan 6, 1967. Ploughed: Jan 11. Basal NPK compound applied: Apr 4. Plots rotary cultivated, potatoes machine planted on OE and CE plots: Apr 20. Weedkiller applied, potatoes dibbled in on OL plots: May 18. Fungicide applied: Aug 3. Sprayed with undiluted BOV at 15 gals: Aug 23. Lifted: Aug 29. Variety: Majestic. Previous crops: Spring wheat 1965, fallow 1966.

Great Hill Bottom I (W): Ploughed: Aug 5, 1966. Ground chalk applied at 40 cwt: Jan 9, 1967. Deep-tine cultivated: Mar 20. Basal NPK compound applied: Mar 20. Rotary cultivated: Apr 20. Potatoes machine planted on OE and CE plots: Apr 21. Weedkiller applied: May 12. Potatoes dibbled in on OL plots: May 18. Sprayed with fentin acetate plus maneb: July 3, July 26, Aug 21. Sprayed with mancozeb: Aug 8. Lifted: Aug 25. Variety: Majestic. Previous crops: Sugar beet 1965, fallow 1966.

NOTE: Samples were taken in June and July for estimates of the incidence of scab and for the degree of tuberisation.

Standard errors per plot. Total tubers:

Pastures (R): 1.117 or 6.4% (6 d.f.)

Great Hill Bottom I (W): 0.771 or 5.8% (6 d.f.)

67/Da/9.2

SUMMARY OF RESULTS

TOTAL TUBERS

OE	OL	CE	Mean
PASTURES (R)			
(±0.559)			
17.93	15.85	18.64	17.47
GREAT HILL BOTTOM I (W)			
(±0.386)			
13.49	11.36	15.12	13.32