

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1967

[Full Table of Content](#)



---

### 67/R/RP11/1/DD/10 Potatoes - Varieties N and Scab

#### Rothamsted Research

Rothamsted Research (1968) *67/R/RP11/1/DD/10 Potatoes - Varieties N and Scab* ; Yields Of The Field Experiments 1967, pp 335 - 339 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/Dd/11.1

POTATOES

(RP 12a/1 and RP 12b/21)

Transmission of scab (*Streptomyces scabies*) - Fosters O & E VI and Highfield O & E III, 1967.

Design (each field): 4 blocks of 5 plots, split into 2.

Area of each sub plot: 0.0014. Area harvested: 0.0013.

Treatments: All combinations of:-

- Whole plots: 1. Seed tuber infection (*Streptomyces scabies*): Severe (SV), moderate (MD), slight (SL), clean (CL), clean seed formalin dipped (FD).  
Sub plots: 2. Chitting: None (O), seed chitted (C).

Basal applications: 10 cwt (13:13:20). Weedkiller: Linuron at 0.75 lb plus paraquat at 0.5 lb ion in 37 gals. Fungicide: Mancozeb at 1.2 lb in 30 gals on three occasions.

Cultivations, etc.: Ploughed: Sept 12, 1966. Basal NPK compound applied: Apr 5, 1967. Plots rotary cultivated, potatoes machine-planted: Apr 26. Weedkiller applied: May 17. Fungicide applied: June 30, July 21, Aug 3. Sprayed with undiluted BOV at 15 gals: Aug 23. Lifted: Sept 22. Variety: Majestic. Previous crops: Fosters - Clover 1965 and 1966, Highfield: Potatoes 1965, barley 1966.

NOTE: Samples were taken during June and July for estimates of the incidence of scab and for the degree of tuberisation.

Standard errors per plot. Total tubers:

Fosters O & E VI. Whole plot: 0.931 or 6.0% (12 d.f.)  
Sub plot: 0.966 or 6.3% (15 d.f.)  
Highfield O & E III. Whole plot: 1.125 or 7.0% (12 d.f.)  
Sub plot: 1.639 or 10.1% (15 d.f.)

67/Da/11.2

SUMMARY OF RESULTS

TOTAL TUBERS

	SV	MD	SL	CL	FD	Mean
FOSTERS O & E VI						
(1) and (2)						(±0.216)
O	14.42	16.59	15.00	16.17	15.67	15.57
C	13.92	17.17	15.25	15.25	14.58	15.23
Mean (±0.465)	14.17	16.88	15.13	15.71	15.13	15.40

HIGHFIELD O & E III

(1) and (2)						(±0.366)
O	14.42	14.92	14.67	16.67	16.08	15.35
C	16.50	18.59	17.75	15.08	17.17	17.02
Mean (±0.563)	15.46	16.75	16.21	15.88	16.63	16.18

Foster O + E VI (1) (±0.577) For use in horizontal and diagonal comparisons  
 (2) (±0.483) For use in vertical and interaction comparisons

Highfield O & E III (1) (±0.808) For use in horizontal and diagonal comparisons  
 (2) (±0.819) For use in vertical and interaction comparisons

67/De/1.1

### SUGAR BEET

Effect of soil compaction on growth of sugar beet, Saxmundham Grove Plot, 1967, the first year.

Design: 4 randomised blocks of 3 plots, split into 4.

Area of each sub-plot: 0.0145. Area harvested: 0.0015.

Treatments: All combinations of:-

- Plots 1. Soil preparation: Soil worked down in winter and allowed to compact (A), soil worked down in spring and compressed (B), soil kept as open as possible when working seedbed (C).
- Sub-plots 2. Fertilisers: 0.6 cwt (N1), 1.2 cwt (N2), 1.8 cwt (N3) 1.2 cwt N plus, 1.6 cwt P2O5 applied as triple superphosphate, half in winter, half in spring (N2P). All N applied in seedbed as 'Nitro-Chalk'.

Basal applications: 0.8 cwt P2O5, 1.0 cwt K2O as triple superphosphate and muriate of potash. Insecticide: Demeton-s-methyl at 3 oz in 37 gals on 2 occasions.

Cultivations, etc.: Ploughed: Nov 29 - Dec 16, 1966. A plots rotary cultivated twice, half phosphate dressing applied to NP sub-plots: Feb 15, 1967. B and C plots worked 3 times with a heavy spring-tine cultivator: Mar 29. All plots harrowed four times with light drag-harrows (twice in each direction), basal PK, 'Nitro-Chalk' and second half phosphate dressing (NP sub-plots) applied, seed drilled at 7 lb. B plots compressed by rolling four times with Cambridge roller: Mar 30. Singled: May 17. Insecticide applied: June 12 and July 7. Lifted: Oct 4. Variety: Sharp's Klein E. Previous crops: Barley 1965 and 1966.

Standard errors per plot.

Roots (washed):	Whole plot:	1.083	or	7.0%	(6 d.f.)
	Sub-plot:	2.053	or	13.3%	(27 d.f.)
Total sugar:	Whole plot:	4.02	or	7.7%	(6 d.f.)
	Sub-plot:	7.37	or	14.1%	(27 d.f.)

67/De/1.2

SUMMARY OF RESULTS

	N1	N2	N3	N2P	Mean
ROOTS (WASHED)					
(1) and (2)					
A	10.66	14.98	17.18	15.20	14.50
B	13.86	14.76	17.79	14.87	15.32
C	13.79	17.03	16.03	19.27	16.53
Mean ( $\pm 0.593$ )	12.77	15.59	17.00	16.45	15.45
SUGAR %					
A	16.9	17.1	16.7	17.3	17.0
B	17.0	16.7	16.4	16.8	16.7
C	17.3	17.2	16.7	16.5	16.9
Mean	17.1	17.0	16.6	16.9	16.9

- (1) ( $\pm 1.041$ ) For use in vertical and diagonal comparisons  
 (2) ( $\pm 1.027$ ) For use in horizontal and interaction comparisons

57/De/1.3

	N1	N2	N3	N2P	Mean
TOTAL SUGAR					
(1) and (2)					
					(±2.01)
A	36.1	51.4	57.5	52.9	49.5
B	47.2	49.4	58.3	50.0	51.2
C	47.8	58.6	53.4	63.6	55.8
Mean (±2.13)	43.7	53.1	56.4	55.5	52.2

PLANT NUMBER					
A	22.3	25.0	24.8	26.6	24.7
B	27.4	27.9	25.5	24.2	26.3
C	26.9	29.8	26.5	31.3	28.6
Mean	25.5	27.6	25.6	27.4	26.5

- (1) (±3.77) For use in vertical and diagonal comparisons  
 (2) (±3.69) For use in horizontal and interaction comparisons