

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/S/DE/1 Sugar Beet - Soil Compaction

Rothamsted Research

Rothamsted Research (1968) *67/S/DE/1 Sugar Beet - Soil Compaction* ; Yields Of The Field Experiments 1967, pp 337 - 338 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/De/1.1

SUGAR BEET

Effect of soil compaction on growth of sugar beet, Saxmundham Grove Plot, 1967, the first year.

Design: 4 randomised blocks of 3 plots, split into 4.

Area of each sub-plot: 0.0145. Area harvested: 0.0015.

Treatments: All combinations of:-

- Plots 1. Soil preparation: Soil worked down in winter and allowed to compact (A), soil worked down in spring and compressed (B), soil kept as open as possible when working seedbed (C).
- Sub-plots 2. Fertilisers: 0.6 cwt (N1), 1.2 cwt (N2), 1.8 cwt (N3) 1.2 cwt N plus, 1.6 cwt P2O5 applied as triple superphosphate, half in winter, half in spring (N2P). All N applied in seedbed as 'Nitro-Chalk'.

Basal applications: 0.8 cwt P2O5, 1.0 cwt K2O as triple superphosphate and muriate of potash. Insecticide: Demeton-s-methyl at 3 oz in 37 gals on 2 occasions.

Cultivations, etc.: Ploughed: Nov 29 - Dec 16, 1966. A plots rotary cultivated twice, half phosphate dressing applied to NP sub-plots: Feb 15, 1967. B and C plots worked 3 times with a heavy spring-tine cultivator: Mar 29. All plots harrowed four times with light drag-harrows (twice in each direction), basal PK, 'Nitro-Chalk' and second half phosphate dressing (NP sub-plots) applied, seed drilled at 7 lb. B plots compressed by rolling four times with Cambridge roller: Mar 30. Singled: May 17. Insecticide applied: June 12 and July 7. Lifted: Oct 4. Variety: Sharp's Klein E. Previous crops: Barley 1965 and 1966.

Standard errors per plot.

Roots (washed):	Whole plot:	1.083	or	7.0%	(6 d.f.)
	Sub-plot:	2.053	or	13.3%	(27 d.f.)
Total sugar:	Whole plot:	4.02	or	7.7%	(6 d.f.)
	Sub-plot:	7.37	or	14.1%	(27 d.f.)

67/De/1.2

SUMMARY OF RESULTS

	N1	N2	N3	N2P	Mean
ROOTS (WASHED)					
(1) and (2)					
A	10.66	14.98	17.18	15.20	14.50
B	13.86	14.76	17.79	14.87	15.32
C	13.79	17.03	16.03	19.27	16.53
Mean (± 0.593)	12.77	15.59	17.00	16.45	15.45

	SUGAR %				
A	16.9	17.1	16.7	17.3	17.0
B	17.0	16.7	16.4	16.8	16.7
C	17.3	17.2	16.7	16.5	16.9
Mean	17.1	17.0	16.6	16.9	16.9

- (1) (± 1.041) For use in vertical and diagonal comparisons
 (2) (± 1.027) For use in horizontal and interaction comparisons