

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1967

[Full Table of Content](#)



---

### 67/R/RG101/DF/1 Grass - Anhydrous Ammonia

#### Rothamsted Research

Rothamsted Research (1968) *67/R/RG101/DF/1 Grass - Anhydrous Ammonia* ; Yields Of The Field Experiments 1967, pp 341 - 343 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/Df/1.1

GRASS

(RG 101)

Anhydrous and aqueous ammonia - Bones Close 1967.

Design: 3 randomised blocks of 26 plots.

Area of each plot: 0.0092. Area harvested: 0.0054.

Treatments: None (NO) - 2 plots per block, and all combinations of:-

1. Nitrogen fertiliser and time of application:

Applied in autumn:

Injected anhydrous ammonia (Nov 15)	IAA
Injected aqueous ammonia (Nov 7)	IQA

Applied in spring:

Injected anhydrous ammonia (Mar 7)	IAS
Injected aqueous ammonia (Mar 8)	IQS

Broadcast 'Nitro-Chalk':

Applied in 3 equal dressings	BD
Applied as single dressing	BS

2. N: 1.0 (N1), 2.0 (N2), 3.0 (N3), 4.0 (N4) cwt  
(total for the season).

Basal application: 900 lb (0:14:28) in winter.

Cultivations, etc.: Basal PK compound applied: Nov 22, 1966. 'Nitro-Chalk' applied (treatment BS and first dressing of BD):  
Mar 22, 1967. Cut three times: May 31, July 20, Oct 11.  
'Nitro-Chalk' applied after first two cuts for BD treatment.

NOTE: Soil samples were taken to determine extent of ammonia diffusion and grass samples for dry matter and percentage of N, P and K.

Standard errors per plot. Grass dry matter

1st cut:	4.46 or 8.8% (46 d.f.)
2nd cut:	2.09 or 11.3% (46 d.f.)
3rd cut:	1.98 or 18.9% (46 d.f.)
Total of 3 cuts:	5.90 or 7.4% (46 d.f.)



67/Df/1.2

SUMMARY OF RESULTS

	IAA	IQA	IAS	IQS	BD	BS	Mean
1ST CUT							
(±2.57)							(±1.05)
N1	47.4	53.6	49.8	45.6	43.7	51.0	48.5
N2	51.2	53.3	47.6	49.2	52.2	46.0	49.9
N3	55.3	54.9	51.1	48.5	48.3	48.4	51.1
N4	57.2	56.6	47.3	48.6	51.1	49.9	51.8
Mean (±1.29)	52.8	54.6	48.9	48.0	48.8	48.8	50.3

NO: 37.2 (±1.82)  
 General mean: 49.3  
 Mean D.M. %: 15.4

2ND CUT							
(±1.20)							(±0.49)
N1	8.2	9.9	16.4	12.3	15.8	12.5	12.5
N2	10.3	21.3	19.6	18.1	23.5	18.0	18.5
N3	16.7	22.3	18.5	22.8	24.6	20.2	20.8
N4	16.5	23.9	20.6	23.8	22.9	25.7	22.2
Mean (±0.60)	12.9	19.4	18.8	19.3	21.7	19.1	18.5

NO: 6.3 (±0.84)  
 General mean: 17.6  
 Mean D.M. %: 24.4

67/Df/1.3

	IAA	IQA	IAS	IQS	BD	BS	Mean
3RD CUT							
(±1.14)							
N1	2.5	2.6	4.6	3.3	10.4	2.9	4.4
N2	3.0	6.7	7.7	5.4	17.5	5.2	7.6
N3	5.8	12.0	11.2	13.5	20.5	9.8	12.1
N4	7.5	18.8	14.0	22.6	22.8	21.2	17.8
Mean (±0.57)	4.7	10.0	9.3	11.2	17.8	9.8	10.5

NO: 2.3 (±0.81)  
 General mean: 58.1  
 Mean D.M. %: 17.7

TOTAL OF 3 CUTS

	(±3.41)						(±1.39)
N1	58.1	66.2	70.7	61.3	69.9	66.5	65.4
N2	64.4	81.4	74.9	72.8	93.1	69.2	76.0
N3	77.8	89.3	80.7	84.8	93.3	78.4	84.1
N4	81.2	99.3	81.8	95.1	96.8	96.8	91.8
Mean (±1.70)	70.4	84.0	77.0	78.5	88.3	77.7	79.3

NO: 45.8 (±2.41)  
 General mean: 76.7  
 Mean D.M. %: 19.2