

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1967

[Full Table of Content](#)



67/R/RG201/DF/2 Grass - N and Damage

Rothamsted Research

Rothamsted Research (1968) *67/R/RG201/DF/2 Grass - N and Damage* ; Yields Of The Field Experiments 1967, pp 345 - 349 - DOI: <https://doi.org/10.23637/ERADOC-1-157>

67/Df/2.1

GRASS

(RG 201)

Nitrogen and damage to sward by ammonia-injectors, Bones Close 1967.

Design: 1 randomised block of 18 plots.

Area of each plot: 0.0092. Area harvested: 0.0054.

Treatments: All combinations of:-

1. Mechanical damage by injector: None (0), damage by 'Anhydrous' injector (Ia), by 'Aqua' injector (Iq).
2. Time of damage: In autumn (A), in spring (S).
3. Nitrogen per cut: 0.3 (N1), 0.6 (N2), 1.0 (N3) cwt as 'Nitro-Chalk' broadcast by hand.

Basal application: 900 lb (0:14:28) applied in winter.

Cultivations, etc.: Autumn injector slits made - 'Aqua': Nov 7, 1966, 'Anhydrous': Nov 14. Basal PK compound applied: Nov 22. Spring injector slits made - 'Anhydrous': Mar 7, 1967, 'Aqua': Mar 8. 'Nitro-Chalk' applied: Mar 22. Cut three times: June 1, July 20, Oct 12. 'Nitro-Chalk' applied after first two cuts.

NOTE: Grass samples were taken for dry matter and percentage of N.

Standard errors per plot (estimated from the 3-factor interaction).

Dry matter:

1st cut:	2.42 or 5.0% (4 d.f.)
2nd cut:	3.26 or 14.4% (4 d.f.)
3rd cut:	0.83 or 5.7% (4 d.f.)
Total of 3 cuts:	3.89 or 4.5% (4 d.f.)

67/Df/2.2

SUMMARY OF RESULTS

DRY MATTER

	IA	IQ	N1	N2	N3	Mean
	(±1.40)		1ST CUT			(±0.99)
A	50.0	53.3	50.7	46.3	58.0	51.7
S	48.3	44.6	43.7	47.7	47.9	46.4
				(±1.71)		(±0.99)
		O	43.8	47.5	54.0	48.4
		IA	50.4	46.6	50.4	49.1
		IQ	43.9	47.4	55.5	48.9
Mean (±0.99)			46.1	47.2	53.3	48.8

Mean D.M. %: 15.8

67/Df/2.3

		DRY MATTER					
		IA	IQ	N1	N2	N3	Mean
		2ND CUT					
		(±1.88)			(±2.31)		(±1.33)
A		23.2	22.3	18.2	24.2	25.8	22.8
S		22.4	22.1	16.9	24.0	25.8	22.3
					(±2.31)		(±1.33)
			0	19.3	23.2	25.8	22.8
			IA	18.0	23.8	26.5	22.8
			IQ	17.1	24.4	25.2	22.2
Mean (±1.33)				18.2	23.8	25.8	22.6

Mean D.M. %: 24.0

67/Df/2.4

		DRY MATTER				
	IA	IQ	N1	N2	N3	Mean
		3RD CUT				
	(±0.48)			(±0.59)		(±0.34)
A	15.0	13.7	9.4	15.6	18.1	14.3
S	14.3	15.7	10.1	16.4	18.6	15.0
				(±0.59)		(±0.34)
		O	9.5	15.6	18.2	14.4
		IA	9.9	15.7	18.3	14.6
		IQ	9.5	16.3	18.4	14.7
Mean (±0.34)			9.6	15.9	18.3	14.6

Mean D.M.%: 18.8

67/Df/2.5

		DRY MATTER				
	IA	IQ	N1	N2	N3	Mean
TOTAL OF 3 CUTS						
	(±2.25)			(±2.75)		(±1.59)
A	88.2	89.3	78.3	86.1	101.9	88.8
S	85.0	82.5	70.7	88.1	92.4	83.7
				(±2.75)		(±1.59)
		0	72.6	86.4	98.0	85.7
		IA	78.4	86.1	95.3	86.6
		IQ	70.6	88.1	99.1	85.9
Mean (±1.59)			73.9	86.9	97.4	86.0

Mean D.M. %: 19.5