

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/B/6 Cultivation - Weedkiller Rotation - Beans, Wheat, Potatoes, Barley

Rothamsted Research

Rothamsted Research (1970) *68/B/6 Cultivation - Weedkiller Rotation - Beans, Wheat, Potatoes, Barley* ; Numerical Results Of The Field Experiments 1968, pp 103 - 109 - DOI:
<https://doi.org/10.23637/ERADOC-1-58>

68/B/6.1

CULTIVATION - WEEDKILLER ROTATION

(CW)

Great Harpenden I 1968 - the 8th year

A comparison of weed-control by various cultivation methods and by pre-emergence weedkillers.

For previous history, rotations, treatments etc., see 'Results' 61/B/10, 62/B/10, 63/B/10, 64/B/9, 65/B/8, 66/B/7 and 'Details' 1967.

A comparison is now made of none v paraquat (G) applied to stubbles after beans, wheat and barley. This is made on half-plots, in all combinations with the test of O v H (post-emergence weedkiller to wheat and barley). The interaction (O v G) x (O v H) is confounded with blocks. Both tests are cumulative, except that in certain years H has been applied to all sub-plots. The first application of paraquat was in autumn 1967, at 0.75 lb ion in 32 gals.

NOTE: As in 1967 A and B plots after potatoes before barley received spring-tine cultivation and no other cultivations.

Area harvested: Beans - 0.0121, wheat, potatoes and barley - 0.0107.

Basal dressing to winter wheat: 2.5 cwt (6:15:15) combine drilled and 3 cwt 'Nitro-Chalk' in spring. All other crops as 1967.

Ground chalk: Spring beans - 23 cwt, winter wheat and potatoes - 46 cwt, barley - 92 cwt.

Cultivations, etc.:-

Spring beans: Paraquat applied to G sub-plots: 12 Oct, 1967. Ground chalk applied: 23 Nov. T plots deep-tine cultivated twice and B plots once: 24 Nov. P and C plots ploughed: 27 Nov. R plots rotary cultivated, depth 6 ins: 6 Dec. P,R,T, B and C plots spring-tine cultivated: 1 Mar, 1968. P,R, T and C plots spring-tine cultivated: 4 Mar. A plots rotary cultivated, seed drilled at 200 lb: 5 Mar. S plots sprayed: 8 Mar. M and C plots tractor-hoed three times: 26 Apr, 20 May and 31 May. Combine harvested: 11 Sept. Variety: Maris Bead.

Winter wheat: Paraquat applied to G sub-plots: 12 Oct, 1967. Ground chalk applied: 23 Nov. T plots deep-tine cultivated twice: 24 Nov. P, C and A plots ploughed: 27 Nov. P,T,A, B and C plots spring-tine cultivated twice, R plots rotary

68/B/6.2

cultivated, seed drilled at 190 lb: 5 Dec. All plots harrowed: 25 Mar, 1968. 'Nitro-Chalk' applied: 12 Apr. H sub-plots and B plots sprayed with mecoprop at 42 oz and 2,4-D at 10.5 oz in 33 gals: 4 May. Combine harvested: 25 Aug. Variety: Cappelle.

Potatoes: Paraquat applied to G sub-plots: 12 Oct, 1967. Ground chalk applied: 23 Nov. T plots deep-tine cultivated twice: 24 Nov. P and C plots ploughed: 27 Nov. R plots rotary cultivated, depth 6 ins: 5 Dec. Basal compound fertiliser applied: 13 Mar, 1968. T plots deep-tine cultivated, P,C and R plots spring-tine cultivated (R plots in error), R plots rotary cultivated, P,C and T plots spring-tine cultivated. A and B plots rotary cultivated, potatoes machine planted: 29 Mar. M and C plots chain harrowed: 26 Apr. S plots sprayed: 3 May. M and C plots grubbed: 20 May, mechanically weeded: 24 May, rotary ridged: 28 May. M, C and Y plots grubbed: 7 June. M, C and Y plots rotary ridged: 12 June. All plots sprayed 3 times with mancozeb at 1.2 lb in 37 gals (the second time including demeton-s-methyl at 3.5 oz): 4 July, 19 July, 5 Aug. Sprayed with undiluted BOV at 15 gals: 30 Aug. Haulm destroyed mechanically 13 Sept. Lifted: 1 Oct.

Barley: All plots spring-tine cultivated twice and sprayed twice with sodium trichloroacetate at 18 lb in 32 gals: 28 Sept and 25 Oct, 1967. All plots spring-tine cultivated: 17 Nov. T plots deep-tine cultivated twice: 24 Nov. Ground chalk applied: 7 Dec. P and C plots ploughed, R plots rotary cultivated, depth 4 - 5 ins: 20 Dec. All plots spring-tine cultivated, seed drilled at 140 lb: 1 Mar, 1968. All plots rolled: 13 Mar. H sub-plots and B plots sprayed with mecoprop at 36 oz and 2,4-D at 9 oz in 33 gals: 4 May. Combine harvested: 21 Aug.

Standard errors per plot.

Spring beans:	Grain, whole plot:	1.43 or 6.9% (8 d.f.)
	sub plot:	0.91 or 4.4% (8 d.f.)
Winter wheat:	Grain, whole plot:	2.40 or 6.5% (8 d.f.)
	sub plot:	1.65 or 4.5% (8 d.f.)
Potatoes:	Total tubers, Whole plot:	1.472 or 8.7% (8 d.f.)
	Sub plot:	0.887 or 5.3% (8 d.f.)
Barley:	Grain, Whole plot:	2.80 or 7.9% (8 d.f.)
	Sub plot:	2.88 or 8.1% (9 d.f.)

68/B/6.3

SUMMARY OF RESULTS

SPRING BEANS

GRAIN

	P	R	T	Mean
Mean (± 0.58)	22.3	21.2	19.9	21.1
M (± 1.01)	22.9	20.4	21.1	21.5 (± 0.58)
S (± 0.71)	22.0	21.6	19.3	21.0 (± 0.41)
	(1) and (2)			
O	22.4	20.4	20.4	21.1
G	22.2	22.0	19.4	21.2

(1) (± 0.64) For use in horizontal and diagonal comparisons only

(2) (± 0.37) For use in vertical and interaction comparisons only

A	AG	B	BG	C	CG
20.4	18.4	16.9	17.2	22.7	22.6

General mean: 20.8

Mean D.M. %: 76.9

68/B/6.4

WHEAT

GRAIN

	P	R	T	Mean
Mean (± 0.98)	35.2	38.1	38.2	37.2
M (± 1.70)	38.0	37.5	38.9	38.1 (± 0.98)
S (± 1.20)	33.9	38.5	37.9	36.7 (± 0.69)
	(1) and (2)			
O	35.2	38.1	38.0	37.1
G	35.2	38.2	38.5	37.3
	(1) and (2)			
O	34.9	38.6	39.1	37.5
H	35.6	37.7	37.3	36.9

(1) (± 1.09) For use in horizontal and diagonal comparisons only
 (2) (± 0.67) For use in vertical and interaction comparisons only

A	AG	AH	AGH	BH	BGH	C	CG
32.9	38.0	34.0	30.4	30.9	35.3	39.8	39.9

General mean: 36.8

Mean D.M. %: 83.2

68/B/6.5

POTATOES: TOTAL TUBERS

	P	R	T	Mean
Mean (± 0.425)	18.19	15.72	16.59	16.84
		(± 0.736)		(± 0.425)
M	17.42	14.91	15.81	16.05
S	18.31	15.71	16.42	16.81
SY	18.83	16.55	17.55	17.65
		(1) and (2)		(± 0.209)
O	18.12	15.47	16.24	16.61
G	18.25	15.98	16.95	17.06

A AG B BG C CG

15.70 17.37 14.86 15.01 19.66 19.01

(1) (± 0.496) For use in horizontal and diagonal comparisons only

(2) (± 0.362) For use in vertical and interaction comparisons only

General mean: 16.86

68/B/6.6

POTATOES: % WARE

	P		R		T	Mean
Mean	98.9	98.9	98.9	98.9	98.9	98.9
M	98.9	98.7	98.8	98.8	98.8	98.8
S	98.8	98.8	98.9	98.9	98.9	98.8
SY	99.0	99.0	99.1	99.1	99.1	99.1
Q	98.9	99.0	98.8	98.8	98.9	98.9
G	98.9	98.7	99.0	99.0	98.9	98.9
A	AG	B	BG	C	CG	
99.0	99.2	98.8	99.2	98.8	99.1	

General Mean: 98.9

68/B/6.7

BARLEY GRAIN				
	P	R	T	Mean
Mean (± 0.81)	36.3	35.0 (± 1.40)	35.8	35.7 (± 0.81)
M	36.5	36.2	35.4	36.0
S	33.9	35.2	36.9	35.3
SY	38.5	33.4	35.0	35.6
		(1) and (2)		(± 0.68)
O	35.2	33.1	33.2	33.9
H	37.3	36.8	38.3	37.5
	A	AH	B	C
	33.4	36.5	34.7	32.2

- (1) (± 1.16) For use in horizontal and diagonal comparisons only
 (2) (± 1.17) For use in vertical and interaction comparisons only

General mean: 35.2

Mean D.M.%: 85.0