

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



### 68/C/5 Cereal Disease Reference Plots - W. Wheat, S. Wheat

#### Rothamsted Research

Rothamsted Research (1970) *68/C/5 Cereal Disease Reference Plots - W. Wheat, S. Wheat ; Numerical Results Of The Field Experiments 1968*, pp 147 - 148 - DOI:

<https://doi.org/10.23637/ERADOC-1-58>

68/C/5.1

CEREAL DISEASE REFERENCE PLOTS

(AQ)

Pennells Piece 1968, the sixth year

For treatments etc., see 'Results' 63/C/10 (WW = Winter wheat, SW = Spring wheat, O = Oats, Be = Spring beans).

Area of each plot: 0.0180. Area harvested: Winter wheat - 0.0115, spring wheat - 0.0115.

Cultivations, etc.: Sprayed plots 1 - 12 (winter wheat blocks) with aminotriazole at 4 lb and ammonium thiocyanate at 3.7 lb in 32 gals: 21 Sept, 1967. Ploughed: 23 Oct. Plots 1 - 12 rotary cultivated: 25 Oct.

Winter wheat: Seed combine drilled at 180 lb: 25 Oct, 1967. 'Nitro-Chalk' applied: 18 Mar, 1968. Sprayed with ioxynil at 7.5 oz and mecoprop at 22.5 oz in 20 gals: 27 Apr. Combine harvested: 24 Aug.

Spring wheat: Seed combine drilled at 175 lb: 6 Mar, 1968. 'Nitro-Chalk' applied: 18 Mar. Sprayed with ioxynil at 7.5 oz and mecoprop at 22.5 oz in 20 gals: 15 May. Combine harvested: 24 Aug.

Oats: Seed combine drilled at 165 lb: 8 Mar, 1968. 'Nitro-Chalk' applied: 18 Mar. Sprayed with ioxynil at 7.5 oz and mecoprop at 22.5 oz in 20 gals: 15 May. Combine harvested: 24 Aug.

Spring beans: Seed placement drilled at 200 lb: 6 Mar, 1968. 1.04 lb phorate applied in granules: 21 June. Combine harvested: 6 Sept.

- NOTES: (1) Yields were taken for winter and spring wheat only (Crop sequences 1, 2, 3 and 6).  
(2) Estimates were made throughout the growing season of the incidence of take-all (*Ophiobolus graminis*) and eyespot (*Cercospora herpotrichoides*).  
(3) For previous year's results see 'Results' 63/C/10, 64/C/9, 65/C/9, 66/C/7 and 67/C/5.

68/c/5.2

SUMMARY OF RESULTS

GRAIN

Crop in	W	W	W	W	
1963	W	W	W	W	
1964	W	W	Be	W	
1965	W	Be	O	W	
1966	Be	O	W	W	
1967	O	W	W	W	Mean
<hr/>					
WINTER WHEAT					
	37.1	34.7	30.1	28.8	32.7
SPRING WHEAT					
	24.2	27.3	25.2	26.1	25.7
Mean D.M. %:	Winter wheat:	84.1			
	Spring wheat:	81.3			