

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Numerical Results of the Field Experiments 1968

[Full Table of Content](#)

INTERNATIONAL DOCUMENTATION SERVICE  
AGRICULTURE  
LONDON AGRICULTURAL BOOKS  
PUBLISHED BY THE  
ROTHAMSTED  
RESEARCH  
UNIT  
ROTHAMSTED, HANTS, RG2 9AT, ENGLAND  
TELEPHONE: 01493 774111 FAX: 01493 774112  
E-MAIL: [eradoc@ Rothamsted.ac.uk](mailto:eradoc@ Rothamsted.ac.uk)  
WWW: <http://www.Rothamsted.ac.uk/eradoc>  
PUBLISHED BY:  
Rothamsted Research

### 68/C/11 Effects of Crop Sequences and Green Manures - Legumes, Barley

#### Rothamsted Research

Rothamsted Research (1970) *68/C/11 Effects of Crop Sequences and Green Manures - Legumes, Barley* ; Numerical Results Of The Field Experiments 1968, pp 171 - 172 - DOI:  
<https://doi.org/10.23637/ERADOC-1-58>

68/C/11.1

## LEGUMES AND BARLEY

(CH)

Effects of crop sequences and green manures - Highfield IV 1968,  
the second year - barley.

Design: 2 randomised blocks of 12 plots, split into 4.

Area of each sub plot: 0.0121. Area harvested: 0.0064.

Treatments: All combinations of:-

Whole plots:

1. Crop sequences (C):

	1	2	3	4	5	6	7
1967	B	B	O	O	Ra	Cl	Be
Undersown	-	T	-	T	-	-	-
1968	B	B	B	B	B	B	B

With nitrogen in 1967: 0.4 (N1), 0.8 (N2) cwt N as 'Nitro-Chalk' to oats and barley, 0.8 (N2), 1.6 (N4) to oilseed rape, none to beans and clover (NO).

Sub plots:

2. Nitrogen to barley 1968: None (T0), 0.35 (T1), 0.70 (T2),  
1.05 (T3) cwt N as 'Nitro-Chalk'.

B = Barley, O = Oats, Ra = Oilseed rape, Cl = Red Clover, Be = Spring beans, T = Trefoil.

Basal applications: 2.5 cwt (0:20:20) combined drilled. Weedkiller: Mecoprop at 36 oz and 2,4-D at 9 oz in 32 gals.

Cultivations, etc.:- Corrective K applied to clover plots at 1.0 cwt K2O as muriate of potash: 22 Nov, 1967. Ploughed: 4 Dec. Seed drilled at 140 lb: 28 Feb, 1968. 'Nitro-Chalk' applied: 15 Mar. Weedkiller applied: 17 May. Combine harvested: 22 Aug. Variety: Maris Badger.

NOTE: (1) Samples were taken on 5 July for estimation of take-all (*Ophiobolus graminis*) and eyespot (*Cercospora herpotrichoides*).

(2) For previous year's results see 'Results' 67/C/27.

Standard errors per plot.

Grain: Whole plot: 1.16 or 3.7% (11 d.f.)

Sub plot: 2.99 or 9.6% (36 d.f.)

68/c/11.2

SUMMARY OF RESULTS

GRAIN

Crop	1967 Under- sown	N	1968				Mean
			NO	N1	N2	N3	
			(1) and (2)				(±0.82)
B	-	1	16.6	28.4	34.7	29.6	27.3
B	-	2	18.8	30.0	35.6	33.5	29.5
B	T	1	33.0	35.4	30.7	27.8	31.7
B	T	2	32.2	35.2	33.1	32.5	33.3
O	-	1	25.0	32.9	34.4	30.1	30.6
O	-	2	23.4	35.2	35.8	29.6	31.0
O	T	1	31.1	36.7	32.3	29.1	32.3
O	T	2	28.4	36.1	30.5	28.7	30.9
Ra	-	2	23.9	36.1	35.8	30.5	31.6
Ra	-	4	29.7	38.6	33.8	27.8	32.5
Cl	-		37.2	29.4	29.2	28.1	30.9
Be	-		28.1	37.0	33.3	29.5	32.0
Mean (±0.61)			27.3	34.2	33.3	29.7	31.1

(1) (±2.01) For use in vertical and diagonal comparisons only

(2) (±2.12) For use in horizontal and interaction comparisons only

Mean D.M. %: 86.6