

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/C/14 Effects of Formalin and Nitrogen - W. Wheat

Rothamsted Research

Rothamsted Research (1970) *68/C/14 Effects of Formalin and Nitrogen - W. Wheat* ; Numerical Results Of The Field Experiments 1968, pp 177 - 180 - DOI:

<https://doi.org/10.23637/ERADOC-1-58>

68/c/14.1

WINTER WHEAT

(EBR, EBS)

Effects of formalin and nitrogen, Pastures (pathogen free 1965), and Little Knott I (pathogen infected 1965) 1968, the fourth year.

Design (each field): 4 x 2 x 2 x 2 x 2 x 2 (quarter replicate) confounded design in 2 blocks of 8 plots split into 2 (Pastures) and in 4 blocks of 8 plots (Little Knott I).

Area of each plot:

Little Knott I: 0.0032. Area harvested: 0.0023.
Pastures (sub plot): 0.0016. Area harvested: 0.0011.

Treatments: All combinations of:-

- Whole plots: 1. N: None (N0), 0.5 (N1), 1.0 (N2), 1.5 (N3) (Pastures) cwt applied as 'Nitro-Chalk' in 1965, 1966 and 1967 and as calcium nitrate in 1968.
2. Formalin applied 1965: (65 O) and (65 F).
Sub plots: 3. Formalin applied 1966: (66 O) and (66 F). (Pastures)
4. Formalin applied 1967: (67 O) and (67 F).
5. Time of application (1967): To wheat stubble (E), after cultivations (L).
6. Formalin applied 1968: None (68 O), sprayed with a 38% solution of formaldehyde at 266 gals in 4220 gals (68 F).

On Little Knott I the treatments were the same, but applied to whole plots only.

Basal applications: 560 lb (0:20:20). Weedkiller: Ioxynil at 9 oz and mecoprop at 27 oz in 50 gals.

Cultivations, etc.:

Pastures: Ploughed: 18 Sept, 1967. Basal PK compound applied: 25 Sept. Formalin applied: 26 Sept. Plots rotary cultivated, seed sown at 180 lb: 24 Oct. Calcium nitrate (first half dressing) applied: 25 Mar, 1968. Weedkiller applied, calcium nitrate (second half dressing) applied: 25 Apr. Combine harvested: 30 Aug. Variety: Cappelle.

68/C/14.2

Little Knott I: Ploughed: 18 Sept, 1967. Basal PK compound applied, formalin applied: 25 Sept. Plots rotary cultivated, seed sown at 180 lb: 24 Oct. Calcium nitrate (first half dressing) applied: 25 Mar, 1968. Weedkiller applied, calcium nitrate (second half dressing) applied: 25 Apr. Combine harvested: 26 Aug. Variety: Cappelle.

- NOTES: (1) Samples of grain and straw were taken at harvest for determination of N.
(2) For previous years' results see 'Results' 65/C/29, 66/C/22, 67/C/18.
(3) Factor 5 has been ignored in the analysis.

Standard errors per plot. Grain:

Pastures (R): Pooled Whole and sub plot: 2.78 or 11.2% (8 d.f.)
Little Knott (R): 4.59 or 14.4% (6 d.f.)

68/c/14.3

SUMMARY OF RESULTS

PASTURES (R)

	NO	N1	N2	N3	Mean
GRAIN					
Mean (± 0.98)	25.9	27.7	23.6	22.1	24.8
		(± 1.39)			(± 0.69)
68 O	28.4	31.0	25.1	23.8	27.1
68 F	23.3	24.3	22.1	20.3	22.5
		(± 1.39)			(± 0.69)
67 O	24.3	27.5	23.1	23.7	24.7
67 F	27.5	27.8	24.1	20.5	25.0
		(± 1.39)			(± 0.69)
66 O	26.2	27.3	23.7	22.2	24.8
66 F	25.6	28.0	23.5	22.0	24.8
		(± 1.39)			(± 0.69)
65 O	27.5	27.2	22.5	21.3	24.6
65 F	24.3	28.2	24.7	22.9	25.0
STRAW					
Mean	30.8	41.9	41.6	37.3	37.9
68 O	32.2	42.5	41.2	36.4	38.1
68 F	29.4	41.2	42.1	38.3	37.7
67 O	33.9	44.0	43.1	35.9	39.2
67 F	27.6	39.7	40.2	38.8	36.6
66 O	31.7	40.7	41.3	36.5	37.6
66 F	29.8	43.0	42.0	38.2	38.2
65 O	31.5	41.8	40.2	34.9	37.1
65 F	30.0	41.9	43.1	39.8	38.7

Mean D.M. %: Grain: 70.2
 Straw: 69.6

68/C/14.4

LITTLE KNOTT I

	NO	N1	N2	N3	Mean
GRAIN					
Mean (± 1.62)	19.3	34.9	37.0	35.8	31.8
		(± 2.29)			(± 1.15)
68 O	18.9	33.1	35.5	37.1	31.1
68 F	19.8	36.8	38.4	34.5	32.4
		(± 2.29)			(± 1.15)
67 O	20.8	34.8	38.6	34.5	32.2
67 F	17.8	35.0	35.3	37.2	31.3
		(± 2.29)			(± 1.15)
66 O	19.2	31.4	35.0	34.8	30.1
66 F	19.4	38.5	38.9	36.8	33.4
		(± 2.29)			(± 1.15)
65 O	20.8	33.2	35.2	34.9	31.0
65 F	17.8	36.7	38.7	36.7	32.5
STRAW					
Mean	20.1	36.6	40.7	39.7	34.3
68 O	20.9	33.5	36.9	40.4	32.9
68 F	19.4	39.6	44.5	39.0	35.6
67 O	23.3	38.1	45.4	39.6	36.6
67 F	16.9	35.1	36.0	39.7	31.9
66 O	20.3	32.2	38.0	39.8	32.6
66 F	20.0	40.9	43.4	39.5	35.9
65 O	22.3	37.0	41.6	36.1	34.2
65 F	18.0	36.2	39.8	43.3	34.3

Mean D.M. %: Grain: 86.2
 Straw: 69.8