

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/C/17 Insecticides and Molluscicides - Grass

Rothamsted Research

Rothamsted Research (1970) *68/C/17 Insecticides and Molluscicides - Grass* ; Numerical Results Of The Field Experiments 1968, pp 187 - 188 - DOI: <https://doi.org/10.23637/ERADOC-1-58>

68/C/25.2

SUMMARY OF RESULTS

DRY MATTER

	O	I	E	G	Mean
STACKYARD (R)					
1ST CUT					
Mean (± 1.12)	21.7	17.9	24.3	29.3	23.3
	(1) and (2)				(± 0.50)
U	17.8	15.2	22.0	28.5	20.9
M	25.5	20.6	26.6	30.2	25.7
	(3) and (4)				(± 0.70)
NO	18.6	11.2	21.7	10.2	15.4
N1	19.6	17.5	23.7	27.1	22.0
N2	24.3	21.1	27.1	37.8	27.6
N3	24.0	21.8	24.7	42.3	28.2

Mean D.M. %: 18.1

(1) (± 1.32) (3) (± 1.65) For use in horizontal and diagonal comparisons only

(2) (± 0.99) (4) (± 1.40) For use in vertical and interaction comparisons only

68/c/25.1

N FIXATION - LUCERNE AND GRASS

(EAT and WEAQ)

Nitrogen fixation by lucerne - Rothamsted (R) Stackyard and Woburn (W) Stackyard C 1968, the second year. For treatments etc. and for the previous year's results see 'Results' 67/c/33.

In 1968 the nitrogen treatments were none (N0), 30 (N1), 60 (N2), 90 (N3) lb N as 'Nitro-Chalk' applied for every cut.

Cultivations, etc.:-

Stackyard (R): PK and first 'Nitro-Chalk' dressings applied: 22 Mar, 1968. Cut 4 times: 15 May, 2 July, 12 Aug and 30 Sept. 'Nitro-Chalk' applied after first 3 cuts.

Stackyard C (W): PK and first 'Nitro-Chalk' dressings applied: 25 Mar, 1968. Cut twice: 20 May, 3 July. 'Nitro-Chalk' applied after each cut.

- NOTES: (1) An error was made in applying the N dressings after the second cut on Stackyard C (W) and the experiment was therefore discontinued.
(2) Soil samples were taken on 2 Feb, 1968 for counts of *Rhizobium leguminosarum*.

Standard errors per plot. Dry matter:

Stackyard (R): 1st cut:	Whole plot:	4.49 or 19.3% (6 d.f.)
	Sub plot:	2.81 or 12.0% (24 d.f.)
2nd cut:	Whole plot:	1.95 or 7.8% (6 d.f.)
	Sub plot:	3.92 or 15.6% (24 d.f.)
3rd cut:	Whole plot:	1.80 or 10.8% (6 d.f.)
	Sub plot:	1.59 or 9.5% (24 d.f.)
4th cut:	Whole plot:	1.26 or 12.1% (6 d.f.)
	Sub plot:	1.46 or 13.9% (24 d.f.)
Total of 4 cuts:	Whole plot:	8.71 or 11.5% (6 d.f.)
	Sub plot:	7.65 or 10.1% (24 d.f.)

Stackyard C (W):

1st cut:	Whole plot:	3.21 or 12.7% (6 d.f.)
	Sub plot:	4.54 or 18.0% (24 d.f.)
2nd cut:	Whole plot:	1.13 or 4.9% (6 d.f.)
	Sub plot:	2.70 or 11.6% (24 d.f.)
Total of 2 cuts:	Whole plot:	3.99 or 8.2% (6 d.f.)
	Sub plot:	6.93 or 14.3% (24 d.f.)