

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



## 68/C/22 Soil Sterilants - W. Wheat

### Rothamsted Research

Rothamsted Research (1970) *68/C/22 Soil Sterilants - W. Wheat* ; Numerical Results Of The Field Experiments 1968, pp 203 - 204 - DOI: <https://doi.org/10.23637/ERADOC-1-58>

68/C/20.1

WINTER WHEAT

(WBW)

Direct seeding, Woburn White Horse Field 1968, the third year.

Design: 4 randomised blocks of 4 plots, split into 2.

Area of each sub plot: 0.0149. Area harvested: 0.0065.

Treatments: All combinations of:-

- Whole plots: 1. Seedbed preparation: Direct seeding after paraquat at 1 lb ion in 33 gals, 21 Sept, 1967 (S). Normal cultivations (M).  
2. Seed dressing: No insecticide, fungicide only, (SO), combined insecticide fungicide (SI).  
Sub plots: 3. Insecticide spray: None (DO), diazinon at 1.6 lb a.i. plus chlordane 1.75 lb a.i. plus DDT at 6.4 lb a.i. plus zinophos at 3.7 lb a.i. in 50 gals (DD).

NOTE: Treatments in 1966, 1967 and 1968 are cumulative.

Basal applications: 45 cwt Magnesian limestone, 5 cwt (0:14:28) broadcast. 0.8 cwt N as 'Nitro-Chalk' top dressed. Weedkiller: Mecoprop at 1.88 lb a.e. in 25 gals.

Cultivations, etc.: 'M' plots ploughed: 20 Sept, 1967. 'S' plots sprayed with paraquat: 21 Sept. Magnesian limestone applied: 12 Oct. PK broadcast: 23 Oct. Insecticide applied, seed drilled at 190 lb, 'S' plots only harrowed: 24 Oct. 'Nitro-Chalk' applied: 22 Apr, 1968. Weedkiller applied: 25 Apr. Combine harvested: 26 Aug. Variety: Cappelle.

NOTES: (1) For the previous years' results see 'Results' 66/C/33 and 67/C/26.

(2) Counts of soil fauna were made in April and plant samples were taken in July for counts of damage to tillers.

Standard errors per plot. Grain:

Whole plot: 3.49 or 12.0% (9 d.f.)  
Sub plot: 6.94 or 23.9% (12 d.f.)

68/c/19.3

SERIES IV

1966 1967 1968	MP MP MP	MP PD MP	PD PD PD	PD MP PD	Mean
TOTAL TUBERS					
(±0.655)					
O	15.71	14.39	3.73	11.78	11.40
C	16.10	14.51	4.21	12.69	11.88
O	15.29	13.34	1.76	10.58	10.24
F	16.52	15.56	6.18	13.89	13.04
Mean (±0.463)	15.90	14.45	3.97	12.23	11.64
% WARE					
O	95.1	95.1	75.4	93.9	89.9
C	94.3	94.1	76.1	94.6	89.8
O	93.7	94.0	66.8	93.1	86.9
F	95.7	95.2	84.8	95.4	92.8
Mean	94.7	94.6	75.8	94.3	89.8

Varieties MP = Maris Piper (resistant)  
 PD = Pentland Dell (susceptible)