Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Numerical Results of the Field Experiments 1968



Full Table of Content

68/C/25 N Fixation by Lucerne - Lucerne and Grass

Rothamsted Research

Rothamsted Research (1970) 68/C/25 N Fixation by Lucerne - Lucerne and Grass; Numerical Results Of The Field Experiments 1968, pp 211 - 220 - **DOI:**

https://doi.org/10.23637/ERADOC-1-58

68/c/17.2

SUMMARY OF RESULTS

DRY MATTER

0	S	В	C	М	P	BCMP	Mean
			1ST CUT				
29.3	28.3	27.6	(±2.81) 27.1	23.5	30.0	23.4	27.0
			2ND CUT				
18.2	20.1	18.8	(±0.75) 17.8	17.9	20.6	16.6	18.6
			3RD CUT				
17.8	19.4	17.8	(±0.58) 17.5	16.4	17.5	16.5	17.5
			4TH CUT				
16.5	16.3	14.7	(±1.00) 15.9	13.7	14.6	14.2	15.1
		TOTAL	OF 2ND,	BRD AND	TH CUTS*		THE BURNEY OF
52.6	55.8	51.3	(±1.45) 51.1	48.1	52.7	47.2	51.3
Mean D	3rd 4th Tot	cut: cut: cut: cut:	20.7 17.4 15.3 17.5 1, 3rd				

^{*} NOTE: The treatments were not applied until after the 1st cut.

68/c/17.1

GRASS

(CS)

Insecticides and molluscicides, Road Piece 1968.

Design: 4 randomised blocks of 7 plots.

Area of each plot: 0.0209. Area harvested: 0.0054.

Treatments: Untreated, no sampling Untreated, sampled	(a) (s)
BHC spray at 6 lb in 200 gals (applied once after first cut)	(B)
Chlorbenside spray at 6 lb in 200 gals (applied 5 times,	(1)
starting after first cut)	(C)
Metaldehyde bait at 5 lb (applied 5 times, starting after first cut)	(M)
Phosdrin spray at 4 oz in 200 gals (applied 5 times,	
starting after first cut)	(P)
Combined treatments	(BCMP)

NOTE: All plots except treatment (0) were sampled as treatment (S).

Basal applications: 3.5 cwt (25:0:16) in spring and 1.75 cwt after each cut except the last.

Cultivations, etc.: Basal NK compound applied: 22 Feb, 1968. Cut 4 times: 28 May, 9 July, 19 Aug, 6 Nov. NK compound applied after first 3 cuts. Insecticides and molluscicides applied:-BHC: 6 June.

Chlorbenside: 11 June, 2 July, 5 Aug, 12 Sept, 3 Oct.

Metaldehyde: 5 June, 1 July, 21 Aug, 4 Sept, 1 Oct.

Phosdrin: 12 June, 2 July, 6 Aug, 4 Sept, 2 Oct.

Previous crop: Grass since 1945 (mainly grazed).

NOTE: Samples were taken for analysis of botanical composition on 27 May, 31 July and 10 Oct. Fauna samples (for estimation of foliage insects, soil insects and slugs) were taken during April, May, July, Sept, Oct and Nov. The concentration of BHC in the soil was determined from samples taken on 10 July, and the soil pH from samples taken on 10 Sept.

Standard errors per plot. Dry matter:

1st cut: 5.62 or 20.8% (18 d.f.) 2nd cut: 1.49 or 8.0% (18 d.f.) 3rd cut: 1.16 or 6.6% (18 d.f.) 4th cut: 1.99 or 13.2% (18 d.f.)

Total of 2nd, 3rd and 4th cuts: 2.89 or 5.6% (18 d.f.)

68/c/16.2

SUMMARY OF RESULTS

	PO	Pl	P4	P6	P24	Mean
			GRAIN	1202		
			(±0.63)			(±0.28)
K1	29.7	35.6	34.3	32.6	34.0	33.2
K ¹ 4	30.2	34.9	31.2	31.7	32.5	32.1
Mean (±0.45) 30.0		35.3	32.7	32.1	33.2	32.7
Mean D.N	1. %: 83.4					
			STRAW			1
Kl	27.6	35.1	37.7	34.9	40.5	35.2
K4	31.4	35.9	39.1	40.0	43.2	37.9
Mean	29.5	35.5	38.4	37.5	41.8	36.5

Mean D.M. %: 89.4

68/c/16.1

BARLEY

(CP/1)

P,K and take-all (Ophiobolus graminis), West Barnfield II 1968, the first year.

Design: 4 randomised blocks of 10 plots.

Area of each plot: 0.0265. Area harvested: 0.0140.

Treatments: All combinations of:-

- 1. Phosphate (in cwt P205 as superphosphate). Applied annually to seedbed: None (P0), 0.3 (P1), 1.2 (P4). Applied in autumn 1967 only, half before ploughing, half after: 1.8 (P6), 7.2 (P24).
- 2. Potash, applied annually to seedbed: 0.3 (K1), 1.2 (K4) cwt K20 as muriate of potash.
- Basal applications: 0.8 cwt N as 'Nitro-Chalk' to seedbed. Weedkillers: Aminotriazole at 4 lb and ammonium thiocyanate at 3.7 lb in 32 gals. Paraquat at 0.75 lb ion in 32 gals.
- Cultivations, etc.: Aminotriazole and ammonium thiocyanate applied: 21 Sept, 1967. Paraquat applied: 6 Oct. Half autumn superphopshate applied: 20 Oct. Ploughed: 26 Oct. Half autumn superphosphate applied: 7 Nov. 'Nitro-Chalk' applied: 29 Feb, 1968. Seedbed P and K applied, seed drilled at 140 lb: 1 Mar. Combine harvested: 21 Aug. Variety: Maris Badger. Previous crops: Grass 1966, winter wheat 1967.
- NOTE: Estimates of eyespot (Cerçosporella herpotrichoides) and takeall (Ophiobolus graminis) were made in spring and summer. Estimates of lodging were also made.

Standard error per plot. Grain: 1.26 or 3.9% (27 d.f.)



68/C/15.3

DRY MATTER

	0	S	Mean
	3RD	CUT	
	(±0.	98)	(±0.69)
L M	11.8 29.6	14.0 28.0	12.9 28.8
Mean (±0.69)	20.7	21.0	20.8

TOTAL OF 3 CUTS

	(±2.	(±1.57)	
L M	54.8 101.6	61.6 98.4	58.2 100.0
Mean (±1.57)	78.2	80.0	79.1

Mean D.M. %: 3rd cut: 18.0 Total of 3 cuts: 17.9

68/C/15.2

SUMMARY OF RESULTS

DRY MATTER

in the second	0	S	Mean
	1ST	CUT	
	(±1.	.41)	(±1.00)
L M	21.8 49.5	25.3 48.2	23.5 48.8
Mean (±1.00)	35.6	36.7	36.2
	2ND	CUT	
ne fit	(±0.	72)	(±0.51)
L M	21.2 22.6	22.3 22.2	21.8
Mean (±0.51)	21.9	22.3	22.1

Mean D.M. %: 1st cut: 15.9 2nd cut: 19.7

68/C/15.1

LUCERNE

(BZ)

Virus control - Long Hoos VI 1968, the third year. For treatments etc. and for previous years' results see 'Results' 66/C/26 and 67/C/20.

Area of each plot: 0.0661. Area harvested: 0.0046.

The dressing applied to M plots after each cut except the last is now 0.5 cwt N as 'Nitro-Chalk' and not NK compound as previously.

Cultivations, etc.: Basal PK compound applied: 14 Dec, 1967. 'Nitro-Chalk' applied to M plots: 12 Mar, 1968. S plots sprayed with demeton-s-methyl at 3.5 oz in 70 gals: 25 Apr, 22 May, 24 June, 13 Aug, 5 Nov. Cut 3 times: 27 May, 5 July, 29 Oct. Muriate of potash applied to L plots and 'Nitro-Chalk' to M plots after first two cuts.

NOTE: Aphid counts were made and the incidences of mosaic virus and of bean leaf roll virus were estimated.

Standard errors per plot. Grass, dry matter: 1st cut: 2.83 or 7.8% (6 d.f.)

2nd cut: 1.44 or 6.5% (6 d.f.) 3rd cut: 1.97 or 9.4% (6 d.f.)

Total of 3 cuts: 4.44 or 5.6% (6 d.f.)

68/c/25.9

DRY MATTER

	0	I	Е	G	Mean
	S	TACKYARD C	(W)		
	TOTAL	OF 1ST AND	2ND CUTS		Ц
Mean (±1.99)	50.8	46.4	51.9	44.6	48.4
		(1) a	nd (2)		(±1.22)
U M	48.1 53.5	42.3 50.5	48.6 55.1	44.2 45.0	45.8 51.0
		(3) a	nd (4)		(±1.73)
NO N1 N2	48.3 56.2 47.9 50.8	41.2 48.3 50.9 45.2	49.7 53.8 55.7 48.2	6.5 40.6 62.8 68.6	36.4 49.7 54.3 53.2

Mean D.M. %: 19.4

^{(1) (±2.64) (3) (±3.60)} For use in horizontal and diagonal comparisons only

^{(2) (±2.45) (4) (±3.46)} For use in vertical and interaction comparisons only

