

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/C/26 Nitrogen Fixation by Lucerne - Lucerne and Grass

Rothamsted Research

Rothamsted Research (1970) *68/C/26 Nitrogen Fixation by Lucerne - Lucerne and Grass* ; Numerical Results Of The Field Experiments 1968, pp 221 - 229 - **DOI:**

<https://doi.org/10.23637/ERADOC-1-58>

68/c/26.1

N FIXATION - LUCERNE AND GRASS

(EBN and WEBO)

Nitrogen fixation by lucerne - Rothamsted (R) Pastures and Woburn (W) Stackyard B, 1968, the first year.

Design: 4 blocks of 4 plots split into 4 for minerals and nitrogen with certain high order interactions confounded with blocks.

Area of each sub-plot: 0.0011. Area harvested: 0.0006.

Treatments: All combinations of:-

- Whole plots: 1. Seed: Lucerne uninoculated (O), lucerne with ineffective inoculum (I), lucerne with effective inoculum (E), Italian ryegrass (G).
- Sub-plots: 2. Minerals: Unmanured (U), 0.6 cwt P₂O₅, 1.2 cwt K₂O as compound fertiliser (0:14:28), plus 4 tons of ground chalk (3 tons on Stackyard B (W)) (M).
3. Nitrogen as 'Nitro-Chalk': None (NO), 30 (N1), 60 (N2), 90 (N3) lb N in the seedbed and after each cut except the last.

Basal applications (Pastures (R) only): Weedkiller: Paraquat at 0.75 lb ion in 37 gals.

Cultivations, etc.:-

Pastures (R): Weedkiller applied: 3 Oct, 1967. Ploughed: 29 Oct. PK, ground chalk and seedbed 'Nitro-Chalk' applied: 28 Mar, 1968. Seed drilled, lucerne at 12 lb, ryegrass at 30 lb: 1 Apr. Cut 3 times: 18 July, 26 Aug, 16 Oct. 'Nitro-Chalk' applied after first 2 cuts. Varieties: Lucerne - Du Puits, ryegrass - S22.

Stackyard B (W): Subsoiled: 27 Sept, 1967. Ploughed: 11 Oct. PK, ground chalk and seedbed 'Nitro-Chalk' applied: 26 Mar, 1968. Seed drilled, lucerne at 12 lb, ryegrass at 30 lb: 27 Mar. Seed re-drilled at same rates because of poor establishment of first drilling: 26 Apr. Cut 3 times: 23 July, 30 Aug, 24 Oct. 'Nitro-Chalk' applied after first 2 cuts. Varieties: Lucerne - Du Puits, ryegrass - S22.

- NOTES: (1) Soil samples were taken before sowing to determine pH and for counts of *Rhizobium leguminosarum*.
- (2) On Stackyard Series B (W) there was flooding of the experiment at the end of April and beginning of May, and also in September.

68/c/26.2

Standard errors per plot. Dry matter:

Rothamsted (Pastures) (R):

1st cut:	Whole plot:	2.34	or	11.0%	(6 d.f.)
	sub plot:	1.88	or	8.9%	(24 d.f.)
2nd cut:	Whole plot:	1.31	or	6.4%	(6 d.f.)
	Sub plot:	1.92	or	9.3%	(24 d.f.)
3rd cut:	Whole plot:	1.43	or	11.8%	(6 d.f.)
	Sub plot:	1.13	or	9.3%	(24 d.f.)
Total of 3 cuts:	Whole plot:	4.09	or	7.6%	(6 d.f.)
	Sub plot:	3.63	or	6.7%	(24 d.f.)

Woburn (Stackyard B) (W):

1st cut:	Whole plot:	2.92	or	15.7%	(6 d.f.)
	Sub plot:	3.04	or	16.4%	(24 d.f.)
2nd cut:	Whole plot:	1.03	or	8.4%	(6 d.f.)
	Sub plot:	2.33	or	18.9%	(24 d.f.)
3rd cut:	Whole plot:	0.90	or	10.2%	(6 d.f.)
	Sub plot:	0.94	or	10.6%	(24 d.f.)
Total of 3 cuts:	Whole plot:	3.11	or	7.8%	(6 d.f.)
	Sub plot:	4.97	or	12.5%	(24 d.f.)

68/c/26.3

SUMMARY OF RESULTS

DRY MATTER

	O	I	E	G	Mean
PASTURES (R)					
1ST CUT					
Mean (± 1.17)	16.7	17.6	21.5	28.7	21.1
		(1) and (2)			(± 0.33)
U	16.7	16.7	18.9	29.3	20.4
M	16.7	18.4	24.2	28.0	21.9
		(3) and (4)			(± 0.47)
NO	9.2	11.2	19.3	16.5	14.0
N1	14.9	15.1	20.8	26.9	19.4
N2	20.1	20.5	22.6	32.8	24.0
N3	22.6	23.5	23.6	38.5	27.1

Mean D.M. %: 20.3

- (1) (± 1.26) (3) (± 1.42) For use in horizontal and diagonal comparisons only
 (2) (± 0.67) (4) (± 0.94) For use in vertical and interaction comparisons only

68/c/26.4

	DRY MATTER				Mean
	O	I	E	G	
	PASTURES (R)				
	2ND CUT				
Mean (± 0.66)	18.9	18.1	22.5	23.2	20.7
	(1) and (2)				(± 0.34)
U	17.9	17.2	21.1	23.8	20.0
M	19.9	19.1	24.0	22.5	21.4
	(3) and (4)				(± 0.48)
N0	10.5	9.2	19.8	9.4	12.2
N1	17.9	15.2	22.6	21.4	19.3
N2	22.3	22.7	24.8	28.0	24.4
N3	24.8	25.3	22.8	33.9	26.7

Mean D.M. %: 16.9

- (1) (± 0.81) (3) (± 1.06) For use in horizontal and diagonal comparisons only
 (2) (± 0.68) (4) (± 0.96) For use in vertical and interaction comparisons only

68/c/26.5

DRY MATTER					
	O	I	E	G	Mean
PASTURES (R)					
3RD CUT					
Mean (± 0.72)	10.3	9.5	11.9	16.8	12.1
	(1) and (2)				(± 0.20)
U	9.4	9.0	10.9	16.9	11.6
M	11.3	9.9	12.9	16.7	12.7
	(3) and (4)				(± 0.28)
NO	6.7	5.3	10.4	7.6	7.5
N1	10.1	8.1	11.5	17.0	11.7
N2	11.2	11.3	13.0	20.7	14.1
N3	13.3	13.2	12.8	21.9	15.3

Mean D.M. %: 15.5

- (1) (± 0.77) (3) (± 0.87) For use in horizontal and diagonal comparisons only
 (2) (± 0.40) (4) (± 0.57) For use in vertical and interaction comparisons only

68/c/26.6

	DRY MATTER				
	O	I	E	G	Mean
	PASTURES (R)				
	TOTAL OF 3 CUTS				
Mean (± 2.04)	45.9	45.2	56.0	68.7	53.9
	(1) and (2)				(± 0.64)
U	44.0	42.9	50.8	70.1	52.0
M	47.9	47.4	61.1	67.2	55.9
	(3) and (4)				(± 0.91)
NO	26.5	25.7	49.4	33.4	33.8
N1	43.0	38.4	54.9	65.3	50.4
N2	53.6	54.5	60.4	81.6	62.5
N3	60.7	62.0	59.1	94.3	69.0

Mean D.M. %: 17.6

- (1) (± 2.24) (3) (± 2.58) For use in horizontal and diagonal comparisons only
 (2) (± 1.28) (4) (± 1.81) For use in vertical and interaction comparisons only

68/c/26.7

	DRY MATTER				Mean
	O	I	E	G	
	STACKYARD B (W)				
	1ST CUT				
Mean (± 1.46)	14.2	13.2	18.8	28.1	18.6
	(1) and (2)				(± 0.54)
U	14.1	11.7	16.0	29.3	17.8
M	14.3	14.7	21.5	27.0	19.4
	(3) and (4)				(± 0.76)
N0	7.9	7.6	14.5	8.8	9.7
N1	14.0	13.6	19.7	20.5	17.0
N2	16.0	13.8	20.3	32.9	20.8
N3	18.8	17.7	20.5	50.3	26.8

Mean D.M. %: 20.4

- (1) (± 1.65) (3) (± 1.97) For use in horizontal and diagonal comparisons only
 (2) (± 1.08) (4) (± 1.52) For use in vertical and interaction comparisons only

68/c/26.8

	DRY MATTER				
	O	I	E	G	Mean
	STACKYARD B (W)				
	2ND CUT				
Mean (± 0.51)	11.2	11.5	13.5	13.0	12.3
	(1) and (2)				(± 0.41)
U	10.3	10.6	12.7	12.4	11.5
M	12.2	12.4	14.4	13.6	13.1
	(3) and (4)				(± 0.58)
NO	5.4	6.0	9.6	2.3	5.8
N1	11.4	10.9	13.9	11.2	11.9
N1	11.9	13.0	15.6	18.4	14.7
N3	16.2	16.0	15.0	20.1	16.8

Mean D.M. %: 16.2

- (1) (± 0.78) (3) (± 1.13) For use in horizontal and diagonal comparisons only
 (2) (± 0.82) (4) (± 1.16) For use in vertical and interaction comparisons only

68/C/26.9

	DRY MATTER				Mean
	O	I	E	G	
	STACKYARD B (W)				
	3RD CUT				
Mean (± 0.45)	6.7	6.5	7.4	14.6	8.8
		(1) and (2)			(± 0.17)
U	6.9	6.4	7.5	14.9	8.9
M	6.5	6.7	7.3	14.3	8.7
		(3) and (4)			(± 0.23)
N0	3.0	3.8	6.2	2.5	3.9
N1	6.0	6.7	7.6	14.2	8.6
N2	8.4	7.7	8.2	19.1	10.8
N3	9.5	8.0	7.6	22.7	11.9

Mean D.M. %: 17.2

(1) (± 0.51) (3) (± 0.61) For use in horizontal and diagonal comparisons only

(2) (± 0.33) (4) (± 0.47) For use in vertical and interaction comparisons only