

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/C/31 Soil Structure 2 - Wheat

Rothamsted Research

Rothamsted Research (1970) *68/C/31 Soil Structure 2 - Wheat* ; Numerical Results Of The Field Experiments 1968, pp 247 - 248 - DOI: <https://doi.org/10.23637/ERADOC-1-58>

68/c/31.1

SOIL STRUCTURE 2

(WEAH)

Effects of peat (annual applications) and subsoiling (1963 only)
Woburn Stackyard Field, plot 6 of the Continuous Barley Site,
red beet 1968, the sixth year.

Design: 4 randomised blocks of 5 plots with treatments to blocks
and strips of sub plots.

Area of each plot: 0.0016. Area harvested: Sub plots without tractor
wheelmarks - 0.0008, with - 0.0005.

Treatments: All combinations of:-

- Blocks: 1. Subsoiling: None (0), subsoiled (S) by hand to a
depth of 20 inches in 1963 (this factor is now
ignored in the analysis).
- Whole plots: 2. Peat (cwt dry matter): None (0), 62.5 applied to top
2 in. of soil in 1965 only (Sb65), 62.5 (DG1), 125 (DG2),
187.5 (DG3) dug in to a depth of 8 in. in 1968.
(DG1 after Sb 1967, DG2 after Dg1 1967, DG3
after Dg2 1967).
- Strips of 5
sub plots: Wheelmarks: 5 rows without wheelmarks (0),
3 rows with wheelmarks (W).

Basal applications: Monoammonium dihydrogen phosphate, potassium
nitrate, ammonium nitrate, and magnesium sulphate to supply
200 lb N, 75 lb P, 250 lb K, 50 lb Mg. All fertilisers
applied to the seedbed before drilling. Insecticide: Dimethoate
at 0.2 pints in 50 gals.

Cultivations, etc.: Peat applied, all plots dug: 29 Jan, 1968. Tractor
wheelmarks made, seedbed fertiliser applied, raked, seed drilled at
30 lb: 15 May. Singled: 20 June. Insecticide applied: 3 July.
Lifted: 9 Sept. Variety: Avon Early.

NOTES: (1) For previous years' results see 'Results' 64/C/20, 65/C/19,
66/C/11, and 67/C/8.

(2) Soil samples for PK analysis were taken in May.

Standard errors per plot.

Marketable roots: Whole plot: 1.506 or 8.4% (12 d.f.)
Sub plot: 1.908 or 10.6% (15 d.f.)

68/c/31.2

SUMMARY OF RESULTS

	O	Sb65	DG1	DG2	DG3	Mean
MARKETABLE ROOTS						
(± 0.954)*						
O	16.81	16.71	18.60	19.27	17.66	17.81
W	18.26	19.31	18.88	18.72	15.88	18.21
Mean (± 0.753)	17.54	18.01	18.74	18.99	16.77	18.01

TOPS FROM MARKETABLE ROOTS						
O	7.14	7.22	7.36	7.75	7.38	7.37
W	7.32	7.69	7.58	7.54	6.68	7.36
Mean	7.23	7.45	7.47	7.64	7.03	7.37

TOTAL PRODUCE						
O	23.95	23.93	25.96	27.02	25.04	25.18
W	25.58	27.00	26.47	26.25	22.56	25.57
Mean	24.77	25.46	26.21	26.64	23.80	25.38

* For use in horizontal and interaction comparisons only