

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/C/32 Levels of N for Winter Wheat After Spring Beans

Rothamsted Research

Rothamsted Research (1970) *68/C/32 Levels of N for Winter Wheat After Spring Beans* ; Numerical Results Of The Field Experiments 1968, pp 249 - 250 - **DOI:**

<https://doi.org/10.23637/ERADOC-1-58>

68/c/32.1

WINTER WHEAT

(WCL)

Levels of N for winter wheat after spring beans - Woburn Lansome III
1968, the second year.

Design: 3 blocks of 4 plots, split into 3.

Area of each sub plot: 0.0072. Area harvested: 0.0019.

Treatments: All combinations of:-

Whole plots: 1. Levels of N to beans 1967: None (R0), 1.0 (R2),
2.0 (R4), 3.0 (R6) cwt N as 'Nitro-Chalk'.

Sub plots: 2. Levels of N 1968: None (N0), 0.5 (N1), 1.0
(N2) cwt N as 'Nitro-Chalk'.

Basal applications: 340 lb (0:14:28) combine drilled. Weedkiller:
Ioxynil/mecoprop (Actril C at 6 pts in 25 gals).

Cultivations, etc.: Sub-soiled both directions: 21 Sept, 1967.
Deep-tine cultivated: 22 Sept. Seed combine drilled at
180 lb: 13 Oct. 'Nitro-Chalk' applied: 17 Apr, 1968.
Weedkiller applied: 23 Apr. Combine harvested: 23 Aug.
Variety: Cappelle.

NOTE: For the previous year's results see 'Results' 67/c/39.

Standard errors per plot. Grain:

Whole plot: 1.92 or 4.8% (6 d.f.)

Sub plot: 2.39 or 6.0% (16 d.f.)

68/c/32.2

SUMMARY OF RESULTS

	GRAIN				Mean
	R0	R2	R4	R6	
	(1) and (2)				(±0.69)
NO	27.9	30.8	31.1	35.2	31.3
N1	42.3	44.8	44.7	47.2	44.7
N2	45.5	43.9	41.8	42.9	43.5
Mean (±1.11)	38.6	39.8	39.2	41.8	39.8

(1) (±1.58) For use in horizontal and diagonal comparisons only

(2) (±1.38) For use in vertical and interaction comparisons only

Mean D.M. %: 79.3