

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/C/33 Nematicides (Formerly Ploughsole D-d) - Sugar Beet

Rothamsted Research

Rothamsted Research (1970) *68/C/33 Nematicides (Formerly Ploughsole D-d) - Sugar Beet ; Numerical Results Of The Field Experiments 1968*, pp 251 - 256 - DOI:
<https://doi.org/10.23637/ERADOC-1-58>

68/C/33.1

NEMATICIDES (FORMERLY PLOUGHSOLE D-D)

(WCM)

The effect of nematicides on Trichodorus and Longidorus in sugar beet - Woburn Butt Furlong, 2nd year.

Design: 2 blocks of 3 plots split into 16.

Area of each sub plot: 0.0045. Area harvested: 0.0034.

Treatments: All combinations of:-

Whole plots: (1) Nitrogen: None (N0), 1.0 (N1), 2.0 cwt N (N2)
as 'Nitro-Chalk'.

Quarter plots: (2) Residues of D-D 1967: None (D0), 65 (D1),
140 (D2), 250 lb D-D (Dichloropropane-dichloro-
propene) (D3).

Strips of three quarter plots (at right angles to quarter plots of factor 2): (3) Fumigants: None (0), 82 lb Telone (dichloropropane) (D) injected in row position before sowing at 6 inch depth. 22.2 lb of 5% granules Lannate (L), 10.6 lb of 10% granules Temik (T). Granules placed in row positions after sowing.

Basal applications: 45 cwt magnesian limestone, 340 lb (0:14:28), 5 cwt salt. Insecticide: Demeton-s-methyl at 3.5 oz in 37 gals.

Cultivations, etc.: Magnesian limestone applied: 7 Nov, 1967. Salt applied, ploughed: 9 Nov. PK applied: 22 Mar, 1968. 'Nitro-Chalk' applied: 25 Mar. Telone applied, seed drilled at 8 lb: 26 Mar. Lannate and Temik applied: 29 Mar. Singled: 20-31 May. Insecticide applied: 17 June. Lifted: 12 Dec. Variety: Klein E.

NOTES: (1) Samples were taken monthly from May to October for nematode counts.

(2) For previous year's results see 'Results' 67/C/40.

Standard errors per plot (pooled):

Roots (washed): 2.087 or 13.2% (47 d.f.)

Total sugar: 6.55 or 13.4% (47 d.f.)

Tops: 2.499 or 18.8% (47 d.f.)

68/c/33.2

SUMMARY OF RESULTS

	O	D	L	T	Mean
ROOTS (WASHED)					
(± 0.738)					
NO	7.57	6.38	7.14	11.10	8.05
N1	18.34	15.43	16.35	21.68	17.95
N2	21.74	20.19	20.17	23.37	21.37
Mean (± 0.426)	15.88	14.00	14.56	18.72	15.79
D0 D1 D2 D3					
(± 0.738)					
NO	5.34	8.43	8.72	9.70	
N1	16.40	18.93	18.57	17.89	
N2	20.44	20.38	22.49	22.17	
(± 0.852)					
O	13.31	16.38	16.70	17.14	
D	12.69	13.91	14.77	14.63	
L	12.68	14.41	15.75	15.39	
T	17.57	18.96	19.15	19.19	
Mean (± 0.426)	14.06	15.92	16.59	16.59	

68/C/33.3

	O	D	L	T	Mean
SUGAR %					
N0	15.7	15.5	15.5	15.8	15.6
N1	15.7	15.8	15.5	15.7	15.7
N2	15.2	15.3	15.2	15.3	15.3
Mean	15.5	15.5	15.4	15.6	15.5
	D0	D1	D2	D3	
N0	15.5	15.7	15.6	15.7	
N1	15.8	15.6	15.7	15.6	
N2	15.3	15.3	15.2	15.2	
O	15.4	15.5	15.6	15.6	
D	15.5	15.5	15.5	15.5	
L	15.4	15.5	15.3	15.3	
T	15.7	15.6	15.6	15.6	
	15.5	15.5	15.5	15.5	

68/c/33.4

	O	D	L	T	Mean
TOTAL SUGAR					
		(±2.31)			(±1.16)
N0	23.8	19.8	22.2	35.1	25.2
N1	57.8	48.6	50.6	68.2	56.3
N2	66.0	61.7	61.2	71.5	65.1
Mean (±1.34)	49.2	43.4	44.7	58.3	48.9

	D0	D1	D2	D3
(±2.31)				
N0	16.7	26.5	27.3	30.5
N1	51.8	59.1	58.5	55.9
N2	62.3	62.2	68.3	67.6
(±2.67)				
O	41.0	50.7	52.0	53.1
D	39.5	42.9	45.9	45.3
L	39.1	44.4	48.1	47.1
T	54.9	59.0	59.4	59.7
Mean (±1.34)	43.6	49.3	51.4	51.3

68/C/33.5

	O	D	L	T	Mean
TOPS					
		(± 0.884)			(± 0.442)
N0	4.83	4.89	5.46	7.67	5.71
N1	13.33	10.88	13.03	17.69	13.73
N2	20.60	19.31	18.75	22.65	20.33
Mean (± 0.510)	12.92	11.70	12.41	16.01	13.26

	D0	D1	D2	D3
(± 0.884)				
N0	3.84	5.72	6.12	7.18
N1	11.77	15.41	13.56	14.19
N2	18.85	19.48	22.29	20.70
(± 1.020)				
O	10.41	13.98	13.27	14.02
D	10.54	11.73	12.66	11.86
L	10.89	12.13	13.58	13.05
T	14.11	16.32	16.45	17.15
Mean (± 0.510)	11.49	13.54	13.99	14.02

