

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)

INTERNATIONAL DOCUMENT SERVICE
AGRICULTURE
LONDON AGRICULTURAL RESEARCH
INSTITUTIONS
OF THE
COMMONWEALTH
COUNCIL
1970
The names of the institutions are given in the list of institutions
at the end of the volume. The names of the authors are given
at the end of each paper. The names of the institutions are given
at the end of each paper. The names of the authors are given
at the end of each paper.

68/C/37 Nematode-resistant Barley

Rothamsted Research

Rothamsted Research (1970) *68/C/37 Nematode-resistant Barley ; Numerical Results Of The Field Experiments 1968*, pp 267 - 268 - DOI: <https://doi.org/10.23637/ERADOC-1-58>

68/C/37.1

NEMATODE-RESISTANT BARLEY

(WCX)

Effect of resistant and susceptible segregates of barley grown in 1967 on land heavily infested with *Heterodera avenae* on the yields of the succeeding susceptible barley - Woburn Butt Close 1968 the 5th year.

For design and previous years' results see 'Results' 64/Da/3, 65/C/30, 66/C/23 and 67/C/42.

Area of each plot: 0.0032. Area harvested: 0.0021.

Treatments: All combinations of:-

1. Nitrogen: 0.4 (N1), 0.8 (N2), 1.2 cwt N (N3) as 'Nitro-Chalk' in the seedbed N1, N2 and N3 being cumulative with N rates 1964 - 67.
2. Formalin 1964: None (64O), Formalin (64F).
3. Formalin 1965: None (65O), Formalin (65F).
4. Formalin 1966: None (66O), Formalin (66F).
5. Varieties 1967: Resistant segregate (R), susceptible segregate (S).

Basal applications: 280 lb (0:20:20) combine drilled. Weedkiller: Ioxynil/mecoprop (Actril C at 5 pts in 25 gals). Magnesian limestone at 22.5 cwt.

Cultivations, etc.:-

Subsoiled: 21 Sept, 1967. Magnesian limestone applied:
18 Oct. Ploughed: 11 Nov. Seed combine drilled at 140 lb:
5 Mar, 1968. 'Nitro-Chalk' applied: 12 Mar. Weedkiller applied:
26 Apr. Combine harvested: 21 Aug. Variety: Maris Badger.

NOTE: Soil samples were taken for nematode counts in June.

Standard error per plot.

Grain: 3.03 or 11.1% (21 d.f.)

68/c/37.2

SUMMARY OF RESULTS

GRAIN

	650	65F	660	66F	R	S	N1	N2	N3	Mean
	(±0.87)		(±0.87)		(±0.87)		(±1.07)			(±0.62)
640	25.9	27.7	27.5	26.1	27.6	26.1	19.4	27.2	34.0	26.8
64F	26.9	29.0	28.2	27.6	29.1	26.8	20.4	28.5	35.0	27.9
			(±0.87)		(±0.87)		(±1.07)			(±0.62)
		650	26.9	25.9	28.4	24.4	18.9	26.9	33.4	26.4
		65F	28.9	27.8	28.3	28.4	20.8	28.7	35.5	28.4
					(±0.87)		(±1.07)			(±0.62)
			660		28.8	26.9	20.0	29.0	34.6	27.9
			66F		27.9	25.9	19.8	26.6	34.3	26.9
							(±1.07)			(±0.62)
					R		21.1	28.9	35.1	28.4
					S		18.7	26.7	33.9	26.4
					Mean		19.9	27.8	34.5	27.4
							(±0.76)			

Mean D.M. %: 83.5