

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/DA/3 Ccc and Eyespot - W. Wheat

Rothamsted Research

Rothamsted Research (1970) *68/DA/3 Ccc and Eyespot - W. Wheat* ; Numerical Results Of The Field Experiments 1968, pp 295 - 296 - DOI: <https://doi.org/10.23637/ERADOC-1-58>

68/Da/3.1

WINTER WHEAT

(RW 201)

Chlormequat and eyespot (*Cercospora herpotrichoides*) - Claycroft
1968.

Design: 4 randomised blocks of 4 plots, split into 2.

Area of each sub plot: 0.0145. Area harvested: 0.0076.

Treatments: All combinations of:-

Whole plots: 1. Chlormequat: * None (0), sprayed** in winter (A),
in spring (S), in winter and in spring (AS).

Sub plots: 2. N: 100 (N1), 200 lb (N2) as 'Nitro-Chalk' in
addition to basal.

* 2-chloroethyltrimethylammonium chloride (CCC).

** At 2 lb in 32 gals on each occasion.

Basal application: 410 lb (6:15:15) combine drilled.

Cultivations, etc.: Ploughed: 13 Sept, 1967. Seed drilled at 180 lb:
5 Oct. Sprayed with chlormequat: 5 Dec and 11 Mar, 1968. 'Nitro-
Chalk' applied: 11 Apr. Combine harvested: 26 Aug. Variety:
Champlain. Previous crops: Barley 1966 and 1967.

NOTE: Samples were taken in spring and summer for estimation of
eyespot (*Cercospora herpotrichoides*).

Standard errors per plot. Grain:

Whole plot: 2.96 or 7.4% (9 d.f.)

Sub plot: 2.59 or 6.5% (12 d.f.)

68/Da/3.2

SUMMARY OF RESULTS

	GRAIN				Mean
	O	A	S	AS	
	(1) and (2)				(±0.65)
N1	40.7	43.0	50.1	42.6	44.1
N2	34.1	35.8	34.0	37.9	35.5
Mean (±1.48)	37.4	39.4	42.1	40.3	39.8

(1) (±1.74) For use in horizontal and diagonal comparisons only

(2) (±1.30) For use in vertical and interaction comparisons only

Mean D.M. %: 83.6