

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Numerical Results of the Field Experiments 1968

[Full Table of Content](#)

INTERNATIONAL DOCUMENT SERVICE  
AGRICULTURE  
LONDON AGRICULTURAL BOOKS  
PUBLISHED BY THE  
ROTHAMSTED  
RESEARCH  
UNIT  
ROTHAMSTED, HANTS, RG2 9AT  
ENGLAND  
PRINTED BY  
HULLMAN LTD.

### 68/DA/4 Spun and Drilled Seed and Cultivations - W. Wheat

#### Rothamsted Research

Rothamsted Research (1970) *68/DA/4 Spun and Drilled Seed and Cultivations - W. Wheat ; Numerical Results Of The Field Experiments 1968*, pp 297 - 298 - DOI:

<https://doi.org/10.23637/ERADOC-1-58>

68/Da/4.1

WINTER WHEAT

(RW 301)

Spun and drilled seed and cultivations - Whittlocks 1968.

Design: 4 randomised blocks of 4 plots, split into 4.

Area of each sub plot: 0.0135. Area harvested: 0.0096.

Treatments:

Spun seed: All combinations of:-

Whole plots: 1. Seed rate: 170 lb (M), 235 lb (H).

Sub plots: 2. Seedbed cultivations: Deep-tine cultivate, spring-tine cultivate twice, sow, harrow (C1).  
Deep-tine cultivate, spring-tine cultivate, sow, spring-tine cultivate, harrow (C2).  
Deep-tine cultivate, spring-tine cultivate twice, harrow, sow, harrow (C3).  
Deep-tine cultivate, sow, spring-tine cultivate twice, harrow (C4).

Drilled seed: All combinations of:-

Whole plot: 1. Seed rate: 150 lb (L), 190 lb (M).

Sub plot: 2. Seedbed cultivations: C1, C3 as above (in duplicate).

Basal applications: 220 lb (0:20:20). 0.84 cwt N as 'Nitro-Chalk' in spring.

Cultivations, etc.: Deep-tine cultivated (with wide points): 19 Oct, 1967. Deep-tine cultivated (with narrow points), basal PK compound applied: 24 Oct. Seed sown: 26 Oct. 'Nitro-Chalk' applied: 11 Apr, 1968. Combine harvested: 26 Aug. Variety: Cappelle. Previous crops: Barley 1966, spring beans 1967.

Standard error per plot. Grain:

Whole plot: 1.45 or 4.5% (6 d.f.)

Sub plot: 2.05 or 6.4% (40 d.f.)

68/Da/4.2

SUMMARY OF RESULTS

GRAIN

SPUN SEED

	C1	C2	C3	C4	Mean
	(1) and (2)				( <del>±0.73</del> ) (±0.73)
M	31.6	31.7	30.8	33.2	31.8
H	32.9	32.9	31.9	32.1	32.5
Mean (±0.73)	32.3	32.3	31.4	32.7	32.2

DRILLED SEED

	C1	C3	Mean
	(3) and (4)		( <del>±0.73</del> ) (±0.73)
L	31.3	31.2	31.3
M	31.6	32.6	32.1
Mean ( <del>±0.51</del> ) (±0.51)	31.5	31.9	31.7

(1) (±1.26) (3) (~~±0.89~~)  
(2) (±1.03) (4) (~~±0.73~~) For use in vertical and diagonal comparisons only  
(±0.73) For use in horizontal and interaction comparisons only

General mean: 31.9  
General mean D.M. %: 84.1