

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/DA/7 Ccc, Row Spacing and N - S. Wheat

Rothamsted Research

Rothamsted Research (1970) *68/DA/7 Ccc, Row Spacing and N - S. Wheat* ; Numerical Results Of The Field Experiments 1968, pp 303 - 304 - DOI: <https://doi.org/10.23637/ERADOC-1-58>

68/Da/7.1

SPRING WHEAT

(RW 451)

Chlormequat, row spacing and N - Geescroft 1968.

Design: 4 randomised blocks of 12 plots.

Area of each plot: 0.0212. Area harvested: 0.0140.

Treatments: All combinations of:-

1. Row spacing: 4 (C), 8 (W) inches.
2. Chlormequat*: None (0), sprayed at 2 lb in 32 gals (S).
3. N: 0.8 (N1), 1.6 (N2), 2.4 (N3) cwt as 'Nitro-Chalk'.

* 2-chloroethyltrimethylammonium chloride (CCC).

Basal applications: 340 lb (0:20:20) broadcast by drill. Weedkillers: Ioxynil/mecoprop (Actril C at 5 pints in 20 gals), barban (Carbyne at 2 pints in 33 gals).

Cultivations, etc.: Ploughed: 15 Sept, 1967. Basal PK applied, seed drilled at 160 lb: 11 Mar, 1968. 'Nitro-Chalk' applied: 21 Mar. Barban applied: 23 Apr. Ioxynil/mecoprop applied, chlormequat applied: 15 May. Combine harvested: 8 Sept. Variety: Kolibri. Previous crops: Fallow 1966, spring oilseed rape 1967.

NOTE: Samples were taken from each plot on 5 June and 3 July for leaf area, shoot height, fresh and dry weight and number of shoots, and for yields on 26 Aug.

Standard error per plot.

Grain: 2.36 or 6.1% (33 d.f.)

68/Da/7.2

SUMMARY OF RESULTS

GRAIN

	O	S	N1	N2	N3	Mean
	(±0.68)			(±0.83)		(±0.48)
C	35.4	41.6	39.1	38.2	38.2	38.5
W	34.0	43.0	37.4	40.7	37.5	38.5
				(±0.83)		(±0.48)
		O	36.4	34.9	32.8	34.7
		S	40.1	43.9	42.9	42.3
		Mean (±0.59)	38.2	39.4	37.9	38.5

Mean D.M. %: 76.3