

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



### 68/DE/4 Effects of Skin-spot (*Oospora Pustulans*) - Potatoes

#### Rothamsted Research

Rothamsted Research (1970) *68/DE/4 Effects of Skin-spot (Oospora Pustulans) - Potatoes ; Numerical Results Of The Field Experiments 1968*, pp 343 - 344 - **DOI:**  
<https://doi.org/10.23637/ERADOC-1-58>

68/De/4.1

## POTATOES

(RP 4/1)

Effects of skin-spot (*Oospora pustulans*) - Long Hoos I and II, 1968.

Design: 6 blocks of 2 plots split into 4.

Area of each sub-plot: 0.0033. Area harvested: 0.0033.

Treatments: All combinations of:-

- Whole plots: 1. Varieties: King Edward (KE), Majestic (M).  
Sub-Plots: 2. Levels of seed infection (*Oospora pustulans*):  
Clean (A), moderately infected (B), severely infected (C), unselected stock (D). B and C were assessed by area covered by skin-spots.

Basal applications: 10 cwt (13:13:20). Weedkiller: Paraquat at 0.38 lb ion plus linuron at 0.75 lb in 36 gals. Fungicide: Mancozeb at 1.2 lb in 37 gals on 3 occasions. Insecticide: Demeton-s-methyl at 3.5 oz applied with mancozeb on one occasion.

Cultivations, etc.: Deep-tine cultivated twice: 2 Oct, 1967.  
Ploughed: 4 - 23 Jan, 1968. Basal NPK applied: 14 Mar.  
Plots rotary cultivated, potatoes machine planted: 9 Apr.  
Weedkiller applied: 4 May. Grubbed and rotary ridged: 17 June. Mancozeb applied: 4 July. Mancozeb and demeton-s-methyl applied: 19 July. Mancozeb applied: 5 Aug. Sprayed with undiluted BOV at 15 gals: 31 Aug. Lifted: 15 Oct.  
Previous crops: Lucerne and cocksfoot 1966, fallow 1967.

NOTE: Emergence counts were made on 20 May and 13 June. After burning off and prior to lifting, count

68/De/4.2

## SUMMARY OF RESULTS

	A	B	C	D	Mean
TOTAL TUBERS					
(1) and (2)					
					( $\pm 0.153$ )
KE	21.46	19.80	17.01	20.08	19.59
M	17.24	17.17	13.18	15.73	15.83
Mean ( $\pm 0.292$ )	19.35	18.49	15.10	17.90	17.71

(1) ( $\pm 0.389$ ) For use in vertical and diagonal comparisons only(2) ( $\pm 0.413$ ) For use in horizontal and interaction comparisons only

	% WARE				
KE	96.9	97.2	97.8	96.9	97.2
M	98.0	98.0	97.9	97.6	97.9
Mean	97.4	97.6	97.8	97.3	97.5