Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Numerical Results of the Field Experiments 1968



Full Table of Content

## **68/DE/5 Rothamsted and Woburn Commercial, Dipped and Healthy Stocks - Potatoes**

#### **Rothamsted Research**

Rothamsted Research (1970) 68/DE/5 Rothamsted and Woburn Commercial, Dipped and Healthy Stocks - Potatoes; Numerical Results Of The Field Experiments 1968, pp 345 - 348 - DOI: https://doi.org/10.23637/ERADOC-1-58

#### POTATOES

### (RP 5/1 and WP 3/1)

Commercial, dipped and healthy stocks - Rothamsted (R) Great Harpenden II and Woburn (W) Lansome III, 1968.

Design: 4 randomised blocks of 6 plots, split into 3.

Area of each sub plot: 0.0036. Area harvested: Rothamsted - 0.0036, Woburn - 0.0036.

Treatments: All combinations of:Whole plots: 1. Seed: Healthy (free from Oospora and Rhizoctonia)
(0), healthy, reinfected with Oospora (CO),
healthy, reinfected with Rhizoctonia (CR), oncegrown, from Rothamsted farm (F), the same, dipped
in fungicide (FD), stock seed (S).

Sub plots: 2. Varieties: King Edward (KE), Majestic (M), Pentland Dell (PD).

Basal applications:
Great Harpenden II (R): 3 tons ground chalk, 7 tons FYM, 8 cwt
(13:13:20). Weedkiller: Paraquat at 0.5 lb ion plus
linuron at 0.75 lb in 40 gals. Fungicide: Mancozeb at 1.2 lb
in 40 gals on 3 occasions.

Lansome III (W): 10 cwt (13:13:20). Weedkiller: Paraquat at 0.38 lb ion plus linuron at 0.5 lb in 50 gals. Fungicide: Mancozeb at 1.2 lb in 38 gals on 3 occasions.

Cultivations, etc.:Great Harpenden II (R): FYM applied: 8 Dec, 1967. Ground chalk applied, plots ploughed: 20 Dec. Basal NPK compound applied: 20 Mar, 1968. Plots rotary cultivated, potatoes planted, plots rotary ridged: 22 Apr. Weedkiller applied: 6 May. Fungicide applied: 5 July, 22 July, 5 Aug. Sprayed with undiluted BOV at 15 gals: 5 Sept. Lifted: 16 Oct. Previous crops: Grass 1959 - 67.

Lansome III (W): Spring-tine cultivated: 11 Mar, 1968. Basal NPK compound applied: 16 Mar. Plots rotary cultivated, potatoes planted: 17 Apr. Grubbed: 25 Apr. Ridged: 26 Apr. Weedkiller

applied: 3 May. Fungicide applied: 12 July, 18 July, 30 July. Sprayed with undiluted BOV at 15 gals: 16 Aug. Lifted: 30 Sept. Previous crops: Spring wheat 1966, fallow 1967.

NOTE: Emergence counts were made on 29 May on Lansome III (W) and on 4 June on Great Harpenden II (R).

Standard errors per plot. Total tubers:

Great Harpenden I (R): Whole plot: 1.107 or 6.0% (15 d.f.)
Sub plot: 1.867 or 10.2% (36 d.f.)

Iansome III (W): Whole plot: 1.157 or 8.2% (15 d.f.)
Sub plot: 1.354 or 9.6% (36 d.f.)

#### SUMMARY OF RESULTS

#### GREAT HARPENDEN II (R)

	0	CO	CR	F	FD	S	Mean
1878,08		T	OTAL TUB	ERS			
- 5.5	(1) and (2)					(±0.381)	
KE	19.11	18.68	17.62	18.28	17.86	20.99	18.76
M PD	19.32 18.85	19.35	17.09 18.46	17.21	18,61	13.23	17.47 18.93
Mean (±0.554)	19.10	19.06	17.73	18.07	18.29	18.07	18.39

<sup>(1) (</sup> $\pm 0.942$ ) For use in horizontal and diagonal comparisons (2) ( $\pm 0.934$ ) For use in vertical and interaction comparisons

	70.		% WARE				1
KE M PD	96.9 98.3 98.5	98.3 98.9 99.0	96.7 98.6 98.9	97 <b>.1</b> 98 <b>.</b> 9 97 <b>.</b> 9	96.1 97.9 97.5	97.7 98.0 99.3	97.1 98.4 98.5
Mean	97.9	98.7	98.0	98.0	97.2	98.4	98.0

#### LANSONE III (W)

	0	CO	CR	F	FD	S	Mean
n aged	18	ŒS.	POTAL TU	BERS	180		1
	(1) and (2)						(±0.276)
KE M PD	13.60 17.15 13.07	14.51 17.49 13.10	12.63 17.51 13.03	12.45 16.60 12.87	12.82 15.56 11.74	13.75 14.01 12.24	13.29 16.39 12.68
Mean (±0.579)	14.60	15.03	14.39	13.98	13.38	13.33	14.12

(1) ( $\pm 0.800$ ) For use in horizontal and diagonal comparisons (2) ( $\pm 0.677$ ) For use in vertical and interaction comparisons

C	***	777
70	WA	1KE

KE	56.5	65.0	55.6	59.6	57.3	73.5	61.2
M	84.1	85.5	86.0	85.9	79.1	87.6	84.7
PD	64.1	70.9	78.5	73.6	57.9	76.8	70.3
Mean	68.3	73.8	73.3	73.0	64.8	79.3	72.1