

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/DE/7 Warm Water Treated Seed - Potatoes

Rothamsted Research

Rothamsted Research (1970) *68/DE/7 Warm Water Treated Seed - Potatoes* ; Numerical Results Of The Field Experiments 1968, pp 353 - 354 - DOI: <https://doi.org/10.23637/ERADOC-1-58>

68/De/7.1

POTATOES

(RP 7/1)

Warm water treated seed - Long Hoos I and II 1968.

Design: 4 randomised blocks of 6 plots.

Area of each plot: 0.0033. Area harvested: 0.0033.

Treatments: None (0) and all combinations of:-

1. Warm water treatment: 45 (L), 50 (M) degrees C.
2. Duration of water treatment: 20 (2), 30 (3) minutes.
Also 55 degrees C (H) for 10 minutes (1).

Basal applications: 10 cwt (13:13:20). Weedkiller: Paraquat at 0.38 lb ion plus linuron at 0.75 lb in 36 gals. Fungicide: Mancozeb at 1.2 lb in 37 gals on 3 occasions. Insecticide: Demeton-s-methyl at 3.5 oz applied with mancozeb on one occasion.

Cultivations, etc. Deep-tine cultivated twice: 2 Oct, 1967.
Ploughed: 4 - 23 Jan, 1968. Basal NPK compound applied: 14 Mar.
Rotary cultivated, potatoes machine planted: 9 Apr.
Grubbed and rotary ridged: 17 June. Mancozeb applied:
4 July. Mancozeb and demeton-s-methyl applied: 19 July.
Mancozeb applied: 5 Aug. Sprayed with undiluted BOV at
15 gals: 31 Aug. Lifted: 15 Oct. Variety: Majestic.
Previous crops: Lucerne and cocksfoot 1966, fallow 1967.

NOTE: Emergence counts were made.

Standard error per plot.

Total tubers: 1.601 or 9.3% (15 d.f.)

68/De/7.2

SUMMARY OF RESULTS

0	L2	L3	M2	M3	H1	Mean
TOTAL TUBERS						
18.49	21.48	19.93	19.33	7.72	16.68	17.27
(±0.800)						
% WARE						
96.3	96.6	96.8	95.8	93.9	96.6	96.0