

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/DE/9 Effects of Aphids - Potatoes

Rothamsted Research

Rothamsted Research (1970) *68/DE/9 Effects of Aphids - Potatoes* ; Numerical Results Of The Field Experiments 1968, pp 357 - 358 - DOI: <https://doi.org/10.23637/ERADOC-1-58>

68/De/9.1

POTATOES

(RP 10/1)

Effects of aphids - Great Harpenden II 1968.

Design: 4 x 4 Latin square.

Area of each plot: 0.0274. Area harvested: 0.0035.

Treatments: All combinations of:-

1. Insecticide applied early: None (0), 1.5 lb phorate as granules applied at planting (E).
2. Insecticide applied late: None (0), sprayed with demeton-s-methyl at 3.5 oz in 37 gals on 5 July (L).

Basal applications: 3 tons ground chalk, 7 tons FYM, 8 cwt (13:13:20). Weedkiller: Paraquat at 0.5 lb ion and linuron at 0.75 lb in 36 gals. Fungicide: Mancozeb at 1.2 lb in 37 gals on 3 occasions.

Cultivations, etc.: FYM applied: 8 Dec, 1967. Chalk applied, plots ploughed: 20 Dec. Basal NPK compound applied: 20 Mar, 1968. Plots rotary cultivated, potatoes machine planted: 11 Apr. Weedkiller applied: 4 May. Rotary ridged: 10 June. Mancozeb applied: 5 July, 19 July, 5 Aug. Sprayed with undiluted BOV at 15 gals: 21 Aug. Lifted: 18 Oct. Variety: King Edward. Previous crops: Grass 1959 - 67.

NOTE: Aphids were counted on leaves removed from potato plants.

Standard error per plot.

Total tubers: 0.648 or 4.2% (6 d.f.)

68/De/9.2

SUMMARY OF RESULTS

	O	L	Mean
TOTAL TUBERS			
(±0.324)			
O	15.53	15.55	15.54
E	15.49	15.68	15.59
Mean (±0.229)	15.51	15.62	15.56
% WARE			
O	95.8	96.3	96.0
E	96.1	95.1	95.6
Mean	96.0	95.7	95.8