Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



# Numerical Results of the Field Experiments 1968



Full Table of Content

# 68/DE/12 Varieties, Nitrogen and Scab (Streptomyces Scabies - Potatoes

#### **Rothamsted Research**

Rothamsted Research (1970) 68/DE/12 Varieties, Nitrogen and Scab (Streptomyces Scabies - Potatoes; Numerical Results Of The Field Experiments 1968, pp 363 - 366 - DOI: https://doi.org/10.23637/ERADOC-1-58

#### POTATOES

# (RP 13/1 and WP 2/1)

Varieties, nitrogen and scab (Streptomyces scabies) - Rothamsted (R) Pastures and Woburn (W) Great Hill Bottom I 1968.

Design: 4 randomised blocks of 3 plots, split into 3.

Area of each sub plot: 0.0120. Area harvested: Pastures (R): 0.0028, Great Hill Bottom I (W): 0.0029.

Treatments: All combinations of:Whole plots: 1. Varieties: King Edward (KE), Majestic (M),
Pentland Dell (PD).
Sub plots: 2. N: None (NO), 1.0 (N1), 2.0 (N2) cwt as 'Nitro-Chalk'.

Basal applications: 6 cwt (0:14:28).

Pastures (R): Weedkillers: Paraquat at 0.75 lb ion in 32 gals.

Paraquat at 0.38 lb ion plus linuron at 0.75 lb in 40 gals.

Fungicide: Mancozeb at 1.2 lb in 37 gals on 3 occasions.

Insecticide: Demeton-s-methyl at 3.5 oz in 37 gals applied with mancozeb on one occasion.

Great Hill Bottom I (W): Weedkillers: Paraquat at 0.5 lb ion in 33 gals. Linuron at 0.5 lb plus paraquat at 0.38 lb ion in 50 gals. Fungicide: Mancozeb at 1.2 lb in 38 gals on 2 occasions.

Cultivations, etc.:

Pastures (R): Paraquat applied: 3 Oct, 1967. Ploughed: 29 Oct.

Basal PK compound applied: 22 Mar, 1968. 'Nitro-Chalk' applied, plots rotary cultivated, potatoes machine planted: 1 Apr.

Paraquat and linuron applied: 3 May. Plots grubbed: 6 June.

Plots rotary ridged: 10 June. Mancozeb applied: 4 July.

Mancozeb and demeton-s-methyl applied: 19 July. Mancozeb applied: 5 Aug. Sprayed with undiluted BOV at 15 gals: 21 Aug. Lifted: 4 Oct. Previous crops: Fallow 1966, barley 1967.

Great Hill Bottom I (W): Deep-tine cultivated: 25 Sept, 1967.

Ploughed: 20 Feb, 1968. Paraquat applied: 3 Mar. Basal PK compound applied: 22 Mar. 'Nitro-Chalk' applied: 3 Apr.

Plots rotary cultivated, potatoes planted: 4 Apr. Linuron and paraquat applied: 2 May. Grubbed and rotary ridged: 15 June. Fungicide applied: 18 and 30 July. Sprayed with undiluted BOV at 15 gals: 15 Aug. Lifted: 4 Oct. Previous crops: Fallow 1966, potatoes 1967.

NOTE: Assessments of surface area of tubers affected by scab (Streptomyces scabies) were made at harvest.

Standard errors per plot. Total tubers:
Pastures (R): Whole plot: 0.285 or 1.7% (6 d.f.)

Sub plot: 1.012 or 6.0% (18 d.f.)

Great Hill Bottom I (W): Whole plot: 0.776 or 8.4% (6 d.f.)

Sub plot: 1.401 or 15.2% (18 d.f.)

https://doi.org/10.23637/ERADOC-1-58

#### SUMMARY OF RESULTS

### PASTURES (R)

	NO	Nl	N2	Mean
		TOTAL TUBE	RS	1 - 1 -
		(1) and (2)		(±0.143)
KE M PD	13.79 13.28 15.15	20.68 14.76 18.13	19.93 15.36 19.86	18.14 14.46 17.71
Mean (±0.292)	14.07	17.86	18.38	16.77

<sup>(1)</sup>  $(\pm 0.437)$  For use in vertical and diagonal comparisons (2)  $(\pm 0.506)$  For use in horizontal and interaction comparisons

		% WARE		
KE M PD	93.2 96.7 98.2	96.4 97.6 99.0	95•7 96•7 98•9	95.1 97.0 98.7
Mean	96.0	97.7	97.1	96.9

## GREAT HILL BOTTOM I (W)

	NO	N1	N2	Mean
	are)d	TOTAL TUBERS	£25	:011
		(1) and (2)		(±0.388)
KE M PD	5.84 7.13 6.36	10.25 10.49 9.13	12.36 10.27 11.20	9.48 9.29 8.90
Mean (±0.404)	6.44	9.96	11.27	9.22

(1)  $(\pm 0.691)$  For use in vertical and diagonal comparisons (2)  $(\pm 0.700)$  For use in horizontal and interaction comparisons

% WARE					
KE M PD	81.7 90.3 89.8	85.8 92.2 92.9	91.4 94.8 94.8	86.3 92.4 92.5	
Mean	87.3	90.3	93.7	90.4	