

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



## 68/DE/13 Woburn the Effect of Chemicals on Common Scab - Potatoes

### Rothamsted Research

Rothamsted Research (1970) *68/DE/13 Woburn the Effect of Chemicals on Common Scab - Potatoes* ; Numerical Results Of The Field Experiments 1968, pp 367 - 368 - DOI:

<https://doi.org/10.23637/ERADOC-1-58>

68/De/13.1

POTATOES

(WP 1/1)

The effect of chemicals on common scab - Woburn Great Hill  
Bottom I 1968.

Design: 4 blocks of 6 plots.

Area of each plot: 0.0064. Area harvested: 0.0021.

Treatments:-

Chemicals: None (O), quintozene at 150 lb (A), captan  
at 50 lb (B), captan at 150 lb (C), drazoxolon  
at 150 lb (D), O,O-diethylphthalimidophosphono-  
thioate at 150 lb (E).

Basal applications: Weedkiller: Paraquat at 0.5 lb ion in  
33 gals. 10 cwt (13:13:20). Weedkiller: Linuron at 0.5 lb  
plus paraquat at 0.37 lb ion in 50 gals. Fungicide: Mancozeb  
at 1.2 lb in 38 gals, applied twice.

Cultivations, etc.: Deep-tine cultivated: 25 Sept, 1967.  
Ploughed: 20 Feb, 1968. Paraquat applied: 3 Mar. NPK  
applied: 22 Mar. Chemical treatments applied, rotary  
cultivated, potatoes planted: 11 Apr. Linuron plus paraquat  
applied: 3 May. Rotary ridged: 15 June. Fungicide applied:  
18 July, 30 July. Sprayed with undiluted BOV at 15 gals:  
16 Aug. Lifted: 4 Oct. Variety: Maris Piper. Previous  
crops: Fallow 1966, potatoes 1967.

NOTE: Scab counts were made.

Standard error per plot.

Total tubers: 0.907 or 7.4% (15 d.f.)

68/De/13.2

SUMMARY OF RESULTS

O	A	B	C	D	E	Mean
TOTAL TUBERS						
(±0.453)						
12.62	14.22	12.75	12.04	11.16	10.36	12.19
% WARE						
90.2	91.9	91.3	89.6	85.5	84.4	88.8