

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Numerical Results of the Field Experiments 1968

[Full Table of Content](#)



68/B/5 Residual Phosphate Rotation - Potatoes, Barley, Swede

Rothamsted Research

Rothamsted Research (1970) *68/B/5 Residual Phosphate Rotation - Potatoes, Barley, Swede* ; Numerical Results Of The Field Experiments 1968, pp 99 - 101 - DOI:
<https://doi.org/10.23637/ERADOC-1-58>

68/B/5.1

RESIDUAL PHOSPHATE ROTATION

(RP)

The long term and residual effects of phosphate fertilisers - Great Field IV and Sawyers I, the 9th year. For treatments and previous years' results see 'Details' 1967 and 'Results' 67/B/6.

Area of each plot:

Great Field IV: 0.0193. Area harvested: Potatoes and barley - 0.0129, swedes - 0.0096.

Sawyers I: 0.0212. Area harvested: Potatoes and barley - 0.0141, swedes - 0.0106.

The basal nitrogen for barley is now 0.8 cwt N on Sawyers and 0.4 cwt N on Great Field IV, applied as 'Nitro-Chalk'.

Cultivations, etc. (both fields):- Ploughed: 20 Nov, 1967.

Potatoes: Fertilisers applied: 28 Mar, 1968. Plots rotary cultivated, potatoes planted: 28 Mar. Sprayed with paraquat at 0.38 lb ion and linuron at 0.75 lb in 36 gals: 3 May. Sprayed 3 times with mancozeb at 1.2 lb in 37 gals: 3 July, 19 July, 5 Aug. The spraying on 19 July included demeton-s-methyl at 3.5 oz. Sprayed with undiluted BOV at 15 gals: 30 Aug. Lifted: 2 Oct.

Barley: Fertilisers applied, seed drilled at 140 lb: 29 Feb, 1968. Combine harvested: 20 Aug.

Swedes: Fertilisers applied: 21 May, 1968. Seed drilled at 1.25 lb: 22 May. Singled: 26 June - 10 July. Lifted: 19 Nov.

Standard errors per plot.

Sawyers I:

Potatoes: 0.899 or 7.0% (11 d.f.)

Barley: 2.29 or 7.5% (11 d.f.)

Swedes: 1.586 or 12.9% (11 d.f.)

68/B/5.2

SUMMARY OF RESULTS

POTATOES

Treatment	TOTAL TUBERS		PERCENTAGE WARE	
	Great Field IV Mean	Sawyers I Mean	Great Field IV Mean	Sawyers I Mean
(±0.636)				
O	11.87	12.27	97.8	97.7
A1	12.91	13.30	97.2	97.8
A2	12.32	13.48	96.2	97.3
A3	13.41	14.46	97.5	96.7
A4	14.25	14.89	96.1	97.2
T1	10.26	9.97	95.4	96.5
T2	13.16	12.20	97.3	98.2
R2	12.12	12.90	96.7	97.5
R3	13.89	13.79	97.0	97.0
R4	15.23	14.81	97.4	97.0
G1	11.42	10.79	96.6	96.6
S1	12.39	11.76	96.6	97.8
Mean	12.77	12.88	96.8	97.3

BARLEY

	GRAIN		STRAW	
(±1.62)				
O	25.4	27.6	31.4	35.3
A1	26.1	31.7	33.0	40.8
A2	25.8	33.5	33.6	50.1
A3	23.1	30.8	43.0	48.4
A4	24.9	31.0	32.5	45.0
T1	25.6	32.0	29.3	37.0
T2	26.8	31.4	32.1	38.2
R2	22.2	27.4	39.8	48.0
R3	26.6	33.4	44.8	44.4
R4	22.4	25.1	59.5	45.0
G1	27.6	32.2	27.5	37.2
S1	25.3	29.8	30.1	39.0
Mean	25.2	30.5	36.4	42.4
Mean D.M. %:	82.0	82.4	86.1	91.7

68/B/5.3

SWEDES, ROOTS

Treat -ment	Great Field IV Mean	Sawyers I Mean
(±1.122)		
O	8.20	3.05
A1	20.09	13.20
A2	21.30	15.15
A3	23.71	15.81
A4	19.77	15.36
T1	14.31	7.32
T2	18.34	10.57
R2	21.21	14.06
R3	22.04	17.76
R4	22.83	17.47
G1	15.37	7.93
S1	11.71	9.70
Mean	18.24	12.28