

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/W/RN/12 - Organic Manuring - Leys & Sugar Beet

Rothamsted Research

Rothamsted Research (1970) *69/W/RN/12 - Organic Manuring - Leys & Sugar Beet* ; Numerical Results Of The Field Experiments 1969, pp 101 - 105 - DOI:

<https://doi.org/10.23637/ERADOC-1-96>

ORGANIC MANURING EXPERIMENT

(69/W/RN/12)

The cumulative effects of organic matter on light land - Woburn Stackyard B 1969, 5th year.

For previous history, rotation, treatments etc., see 'Results' 66/C/31, 67/C/24 and 68/C/18. All plots except those under leys (LC and LN) carried sugar beet.

Area of each sub plot: 0.0156. Area harvested: Leys: 0.0129.
Sugar beet: 0.0034.

Fertilisers applied Autumn 1968 (cwt)

Treatment	P2O5	K2O	MgO
DG	-	-	-
ST	0.4	-	0.15
PT	0.5	1.0	-
GM	0.5	1.0	0.2
FD	1.0	3.0	0.8
FS	0.5	1.0	0.2
LC	0.8	1.0	0.2
LN	0.8	1.0	0.2

Fertilisers applied Spring 1969 (cwt)

Treatment	P2O5	K2O
DG	-	-
ST	0.5	1.5
PT	0.5	1.0
GM	0.7	1.1
FD	-	1.6
FS	0.5	1.0
LC	0.4	1.2
LN	0.4	1.0

No magnesium was required in the Spring.

Nitrogen to sugar beet:

N1, N3, N5, N7. 0.2, 0.6, 1.0, 1.4 cwt N as 'Nitro-Chalk'.

Basal and standard applications: Ground chalk to whole area at 4.5 tons.

Insecticide to sugar beet: Demeton-s-methyl at 3.5 oz in 30 gals.

Cultivations, etc.:

LC and LN plots: P,K, and Mg applied: 8 Nov, 1968. Ground chalk applied: 10, 14 Feb, 1969. P,K applied: 3 Mar. N applied to LN plots: 14 Mar, 3 July. Cut: 25 June, 3 Sept.

Sugar beet: P, K, and Mg applied: 19, 20 Nov, 1968. Peat, straw, FYM applied: 28 - 29 Nov. Ploughed: 29 Nov. Ground chalk applied: 10, 14 Feb. NPK applied: 10 - 11 Apr. Power harrowed, seed drilled at 5 lb: 11 Apr. Singled: 29 May - 3 June. Insecticide applied: 23 June. Lifted: 29 - 30 Oct. Variety: Klein E.

Standard errors per plot. Sugar beet:

Roots (washed), tons:	Whole plot: 1.183 or 8.9% (15 d.f.)
	Sub plot: 1.342 or 10.1% (54 d.f.)
Total sugar, cwt:	Whole plot: 5.31 or 10.2% (15 d.f.)
	Sub plot: 5.50 or 10.6% (54 d.f.)
Tops, tons:	Whole plot: 0.759 or 13.0% (15 d.f.)
	Sub plot: 1.161 or 10.8% (54 d.f.)

SUMMARY OF RESULTS

SUGAR BEET

	N1	N3	N5	N7	Mean
ROOTS (WASHED): TONS					
(1) and (2)					
					(±0.592)
DG	14.68	15.97	16.70	16.47	15.96
ST	9.80	12.37	13.77	14.45	12.60
PT	9.67	11.56	14.57	15.11	12.73
GM	13.34	14.11	14.70	14.44	14.15
FD	9.12	12.34	13.25	13.74	12.11
FS	9.63	12.34	13.51	13.64	12.28
Mean (±0.274)	11.04	13.11	14.42	14.64	13.30

(1) (±0.829) For use in vertical and diagonal comparisons only

(2) (±0.671) For use in horizontal and interaction comparisons only

SUGAR %

DG	20.1	20.0	19.5	19.2	19.7
ST	19.8	20.0	20.0	19.3	19.8
PT	19.9	20.2	19.3	19.5	19.7
GM	19.7	19.7	19.0	18.6	19.3
FD	19.9	19.7	19.3	18.7	19.4
FS	19.6	19.6	19.9	19.0	19.6
Mean	19.9	19.8	19.5	19.0	19.6

SUGAR BEET

	N1	N3	N5	N7	Mean
TOTAL SUGAR: CWT					
	(1) and (2)				(±2.65)
DG	59.0	63.7	65.1	63.2	62.8
ST	38.9	49.5	55.4	55.9	49.9
PT	38.4	46.6	56.4	58.9	50.1
GM	52.7	55.5	56.1	53.7	54.5
FD	36.4	48.9	51.9	52.3	47.4
FS	37.8	48.6	54.2	52.5	48.3
Mean (±1.12)	43.9	52.1	56.5	56.1	52.2

(1) (±3.57) For use in vertical and diagonal comparisons only

(2) (±2.75) For use in horizontal and interaction comparisons only

TOPS: TONS

	(1) and (2)				(±0.380)
DG	4.89	6.55	7.87	8.47	6.95
ST	2.74	4.10	5.69	7.14	4.92
PT	3.04	4.30	6.88	7.61	5.46
GM	5.75	6.88	8.53	9.66	7.71
FD	3.17	4.40	5.72	6.81	5.03
FS	2.94	4.76	5.29	7.08	5.02
Mean (±0.237)	3.76	5.16	6.66	7.79	5.85

(1) (±0.630) For use in vertical and diagonal comparisons only

(2) (±0.580) For use in horizontal and interaction comparisons only

LEY: DRY MATTER

	LC		LN
	1ST CUT		
	26.0		36.0
	2ND CUT		
	4.1		8.5
	TOTAL OF 2 CUTS		
	30.1		44.5

Mean D.M. %: 1st cut: 29.2
 2nd cut: 33.3
 Total of 2 cuts: 31.2