

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/R/CS/12 - Formalin & N (2 Sites) - Winter Wheat

Rothamsted Research

Rothamsted Research (1970) *69/R/CS/12 - Formalin & N (2 Sites) - Winter Wheat* ; Numerical Results Of The Field Experiments 1969, pp 140 - 142 - **DOI:**

<https://doi.org/10.23637/ERADOC-1-96>

WINTER WHEAT

(69/R/CS/12)

Effects of formalin and nitrogen, Pastures (pathogen free 1965) and Little Knott I (pathogen infected 1965) 1969, the fifth year. For design, treatments etc., and previous years' results see 'Results' 65/C/29, 66/C/22, 67/C/18 and 68/C/14.

Area of each plot: 0.0032. Area harvested: Little Knott I: 0.0021.
Pastures: 0.0010.

No formalin treatment was applied in 1969, residual effects only being measured. Nitrogen was applied as calcium nitrate as in 1968 and at the same rates.

Basal applications: Compound (0:20:20) at 280 lb on Pastures and at 168 lb on Little Knott I, combine drilled. Weedkiller: Ioxynil octanoate, bromoxynil octanoate and the iso-octyl ester of dichlorprop ('Oxytril P' at 1 pint in 20 gals).

Cultivations, etc.:

Pastures: Ploughed: 14 Oct, 1968. Seed drilled at 180 lb: 17 Oct. Calcium nitrate applied: 16 Apr, 1969. Weedkiller applied: 1 May. Combine harvested: 1 Sept. Variety: Cappelle.

Little Knott I: Ploughed: 20 Sept, 1968. Seed drilled at 180 lb: 15 Oct. Calcium nitrate applied: 16 Apr, 1969. Weedkiller applied: 1 May. combine harvested: 1 Sept. Variety: Cappelle.

NOTE: Samples were taken for estimation of take-all (*Ophiobolus graminis* and eyespot (*Cercospora herpotrichoides*) in spring and early and late summer.

Standard errors per plot.

Grain, cwt:	Little Knott I:	4.36 or 15.2% (6 d.f.)
	Pastures: Whole plot:	1.74 or 5.0% (5 d.f.)
	Sub plot:	4.24 or 12.2% (3 d.f.)
	Pooled:*	3.24 or 9.3% (8 d.f.)

* Used in calculating S.E.'s of means in Summary.

SUMMARY OF RESULTS

PASTURES (R)

GRAIN: CWT

	NO	N1	N2	N3	Mean
Mean (± 1.15)	18.0	37.2	42.8	41.4	34.9
		(± 1.62)			(± 0.81)
68 O	19.0	38.7	41.3	40.9	35.0
68 F	17.0	35.8	44.2	41.9	34.7
		(± 1.62)			(± 0.81)
67 O	16.8	38.0	42.8	42.9	35.1
67 F	19.2	36.5	42.7	40.0	34.6
		(± 1.62)			(± 0.81)
66 O	17.8	37.0	41.1	41.2	34.3
66 F	18.2	37.5	44.5	41.6	35.4
		(± 1.62)			(± 0.81)
65 O	16.6	35.9	40.1	43.1	33.9
65 F	19.4	38.6	45.5	39.8	35.8

Mean D.M. %: 82.8

LITTLE KNOTT I

GRAIN: CWT

	NO	N1	N2	N3	Mean
Mean (± 1.54)	14.4	30.7	32.8	36.8	28.7
		(± 2.18)			(± 1.09)
68 O	14.4	32.4	37.5	38.9	30.8
68 F	14.5	29.0	28.1	34.7	26.6
		(± 2.18)			(± 1.09)
67 O	13.9	29.8	31.3	33.5	27.1
67 F	15.0	31.6	34.3	40.1	30.3
		(± 2.18)			(± 1.09)
66 O	15.0	27.6	32.3	39.8	28.7
66 F	13.9	33.8	33.3	33.8	28.7
		(± 2.18)			(± 1.09)
65 O	13.3	31.1	33.0	32.6	27.5
65 F	15.6	30.3	32.6	41.0	29.9

Mean D.M. %: 81.6