


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/W/CS/17 - Cereal Cyst Nematode - Barley

Rothamsted Research

Rothamsted Research (1970) *69/W/CS/17 - Cereal Cyst Nematode - Barley* ; Numerical Results Of The Field Experiments 1969, pp 160 - 163 - DOI: <https://doi.org/10.23637/ERADOC-1-96>

CEREAL CYST-NEMATODE

(69/W/CS/17)

The effect of cereal cyst-nematode (*Heterodera avenae*) on the yield of resistant barley - Butt Close 1969, fourth year. For treatments, previous years' results etc., see 'Results' 67/C/41 and 68/C/36.

Design: 6 blocks of 5 plots split into 3.

Area of each sub plot: 0.0031. Area harvested: 0.0020.

Treatments: All combinations of:-

- Blocks: 1. Crop in 1968: Spring wheat, spring barley in three pairs of blocks.
- Whole plots: 2. Oats 1966: Under oats till harvest (O), oats rotary cultivated 26 May, and then bare fallowed (A), oats rotary cultivated 26 May, bare fallowed, injected with D-D at 400 lb, 19 Dec, 1966 (B).
- Sub plots: 3. Cumulative nitrogen to wheat 1967, 1968: 0.6 (N1), 1.2 (N2), 1.8 (N3) cwt N as 'Nitro-Chalk'.

or

Cumulative nitrogen to barley 1967, 1968: 0.4 (N1), 0.8 (N2), 1.2 (N3) cwt N as 'Nitro-Chalk'.

NOTE: In 1968 because of a severe invasion of grain aphid *Sitobion avenae* a test of 0 v. 0.4 lb dimethoate (S) in 30 gals was made on spring wheat, two of the three O plots in each block and the A and B plots were sprayed.

Basal applications: 350 lb (20:10:10) combine drilled. Weedkiller: Ioxynil at 7.5 oz and mecoprop at 22.5 oz in 25 gals.

Cultivations, etc.:- Ploughed: 3 Feb, 1969. Seed combine drilled at 154 lbs: 26 Mar. Weedkiller applied: 14 May. Combine harvested: 7 Aug. Variety: Resistant Barley.

- NOTES: 1. Plant samples were taken for nematode cyst counts on roots 2 June.
2. Soil samples were taken for nematode counts after harvest.

Standard errors per plot. Grain, cwt:

After Spring wheat:	Whole plot:	1.52 or 4.1% (8 d.f.)
	Sub plot:	3.31 or 9.0% (20 d.f.)
After Barley:	Whole plot:	1.86 or 5.6% (8 d.f.)
	Sub plot:	4.32 or 12.9% (20 d.f.)
Pooled*	Whole plot:	1.70 or 4.8% (16 d.f.)
	Sub plot:	3.85 or 11.0% (40 d.f.)

* Used in calculation of SE's of means in Summary.

SUMMARY OF RESULTS

BARLEY, GRAIN

AFTER SPRING WHEAT

	OS	O	AS	BS	Mean
	(1) & (2)		(3) & (4)		(±0.99)
N1	35.8	37.6	37.5	41.7	37.7
N2	33.4	35.2	36.2	37.3	35.1
N3	36.4	36.9	36.5	40.6	37.4
Mean (±0.98)	35.2 (±0.69)	36.5	36.7	39.9	36.7

(1) (±1.46) (3) (±2.06) For use in horizontal and diagonal comparisons only.

(2) (±1.57) (4) (±2.22) For use in vertical and interaction comparisons only.

BARLEY, GRAIN

AFTER BARLEY

	O	A	B	Mean
	(1) & (2)	(3) & (4)		(±0.99)
N1	31.0	35.3	39.0	33.5
N2	33.8	32.9	33.5	33.5
N3	31.7	29.9	41.4	33.3
Mean (±0.98)	32.2 (±0.69)	32.7	37.9	33.4

(1) (±1.46) (3) (±2.06) For use in horizontal and diagonal comparisons only.

(2) (±1.57) (4) (±2.22) For use in vertical and interaction comparisons only.

Pooled mean: 35.1
Pooled mean D.M.%: 85.8