

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



### 69/S/CS/4 - P and K Residues - Spring Wheat After Clover

#### Rothamsted Research

Rothamsted Research (1970) *69/S/CS/4 - P and K Residues - Spring Wheat After Clover* ; Numerical Results Of The Field Experiments 1969, pp 233 - 234 - DOI:

<https://doi.org/10.23637/ERADOC-1-96>

SPRING WHEAT

(69/S/CS/4)

Phosphate and potash, Saxmundham, Victor's Plot, 1969, the third year, spring wheat following clover. For treatments etc. and for previous years' results see 'Results' 67/C/45 and 68/C/41.

Design: 4 randomised blocks of 4 plots, with plots split into 2 for N\*.

Area of each plot: 0.0018. Area harvested: 0.0011.

Treatments: All combinations of:-

- Whole plots: 1. Phosphate: 0.5 cwt (P1), 2.0 cwt (P4) P<sub>2</sub>O<sub>5</sub> as triple superphosphate.
2. Potash: None (K0), 2.0 cwt (K1) K<sub>2</sub>O as muriate of potash.
- Sub plots\*: 3. Nitrogen: 0.5 cwt (N1), 1.0 cwt (N2) as 'Nitro-Chalk'.

Basal applications: Manures: None. Weedkiller: Paraquat at 0.5 lb ion in 25 gals.

Cultivations, etc.: P and K applied: 25 Sept, 1968. Ploughed: 31 Oct - 14 Nov, 1968. Paraquat applied: 17 Apr, 1969. Seed drilled: 18 Apr. N applied: 14 May. Combine harvested: 21 Aug. Variety: Kolibri.

Standard error per plot.

Grain, cwt: 1.23 or 8.3% (8 d.f.)

\* NOTE: The split for N was ignored at harvest.

SUMMARY OF RESULTS

	KO	KI	Mean
	GRAIN: CWT		
	(±0.62)		(±0.44)
P1	15.6	13.8	14.7
P4	15.4	14.5	14.9
Mean (±0.44)	15.5	14.1	14.8
	STRAW: CWT		
P1	30.9	32.0	31.5
P4	29.1	30.4	29.8
Mean	30.0	31.2	30.6

Mean D.M. %: Grain: 78.7  
 Straw: 63.1