

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



### 69/R/W/ 10 - Chemical Control of Take-all -

#### Rothamsted Research

Rothamsted Research (1970) *69/R/W/ 10 - Chemical Control of Take-all -* ; Numerical Results Of The Field Experiments 1969, pp 249 - 250 - DOI: <https://doi.org/10.23637/ERADOC-1-96>

WINTER WHEAT

(69/R/W/10)

Chemical control of take-all (*Ophiobolus graminis*), Highfield Drive 1969.

Design: 4 randomised blocks of 7 plots.

Area of each plot: 0.0050. Area harvested: 0.0026.

Treatments: None (0) and all combinations of:-

1. Fungicide: Benomyl (B), M2452(M), each as 50 % wettable powder, 'Plantvax' (P) as 75% wettable powder.
2. Rates of application: 1 lb(1), 5 lb(2) a.i.

Basal applications: 280 lb (6:15:15) combine drilled. 0.8 cwt N as 'Nitro-Chalk' in spring. Weedkiller: mecoprop at 42 oz and 2,4-D at 10.5 oz in 20 gals.

Cultivations, etc.: Ploughed: 20 Sept, 1968. Seed drilled at 180 lb: 16 Oct. 'Nitro-Chalk' applied: 27 Mar, 1969. Fungicides applied: 10 Apr. Weedkiller applied: 1 May. Combine harvested: 30 Aug. Variety: Cappelle. Previous crops: Barley 1967 and 1968.

NOTE: Samples were taken for estimation of take-all (*Ophiobolus graminis*), eyespot (*Cercospora herpotrichoides*) and sharp eyespot (*Rhizoctonia solani*) in spring and summer.

Standard error per plot.

Grain, cwt: 6.07 or 12.6% (15 d.f.)

SUMMARY OF RESULTS

GRAIN: CWT

	B	M	P	Mean
		(±3.04)		(±1.75)
1	50.2	45.6	46.9	47.6
2	51.2	49.8	45.3	48.8
Mean (±2.15)	50.7	47.7	46.1	48.2

0 46.6 (±3.04)

General Mean: 48.0

Mean D.M.%: 81.9