

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



### 69/R&W/B/1 - Varieties, N & Ethirimol - Barley

#### Rothamsted Research

Rothamsted Research (1970) *69/R&W/B/1 - Varieties, N & Ethirimol - Barley* ; Numerical Results Of The Field Experiments 1969, pp 257 - 259 - DOI: <https://doi.org/10.23637/ERADOC-1-96>

BARLEY

(69/R/B/1 and 69/W/B/1)

Varieties, N and ethirimol, ('Milstem' or 'PP149') Rothamsted (R)  
Pastures and Woburn (W) Lansome II 1969.

Design (each experiment): 4 blocks of 8 plots (randomisation  
restricted), split into three for N.

Area of each sub plot: 0.0080. Area harvested: 0.0054.

Treatments: All combinations of:-

- Whole plots: 1. Varieties: Julia (J), Maris Badger (MB),  
Sultan (S), Zephyr (Z).  
2. Fungicide: None (O), ethirimol as seed  
dressing at 1 lb (F).  
Sub plots: 3. Nitrogen: 0.3 (N1), 0.6 (N2), 0.9 (N3)  
cwt as 'Nitro-Chalk'.

Basal applications: 220 lb (0:20:20) combine drilled.

Weedkillers: Pastures (R):- Paraquat at 0.5 lb ion in 25  
gals. 2,4-D at 8 oz and dichlorprop at 32 oz in 20 gals.  
Lansome II (W):- Ioxynil octanoate, bromoxynil octanoate  
and the iso-octyl ester of dichlorprop ('Oxytril P' at 1 pint  
in 25 gals).

Cultivations, etc.:

Pastures (R): Paraquat applied: 11 Oct, 1968. Ploughed:  
13 Nov. Seed drilled at 140 lb: 27 Mar, 1969. 'Nitro-Chalk'  
applied: 10 Apr. 2,4-D/dichlorprop applied: 13 May. Combine  
harvested: 14 Aug. Previous crops: Potatoes 1967, barley, 1968.  
Lansome II (W): Ploughed: 18 Nov, 1968. Seed drilled at  
140 lb: 28 Mar, 1969. 'Nitro-Chalk' applied: 8 Apr. Weedkiller  
applied: 14 May. Combine harvested: 14 Aug. Previous crops:  
Spring beans 1967 Winter wheat 1968.

NOTE: Samples were taken for assessment of mildew (*Erysiphe  
graminis*) and other foliar diseases.

Standard errors per plot.

Grain, cwt: Pastures (R): Whole plot: 1.74 or 3.6% (21 d.f.)  
Sub plot: 2.04 or 4.2% (48 d.f.)  
Lansome II (W): Whole plot: 1.14 or 2.8% (21 d.f.)  
Sub plot: 1.87 or 4.5% (48 d.f.)

SUMMARY OF RESULTS

GRAIN:CWT

PASTURES (R)

	J	MB	S	Z	Mean
	(±0.87)				(±0.43)
O	49.6	41.4	49.5	48.6	47.3
F	51.5	43.6	48.4	51.2	48.7
	(1) and (2)				(±0.36)
N1	45.9	38.4	43.2	47.5	43.7
N2	52.5	45.5	52.3	51.7	50.5
N3	53.3	43.6	51.4	50.7	49.7
Mean (±0.61)	50.6	42.5	48.9	49.9	48.0

Mean D.M.%: 83.6

- (1) (±0.85) For use in horizontal and diagonal comparisons only  
 (2) (±0.72) For use in vertical and interaction comparisons only

SUMMARY OF RESULTS

GRAIN: CWT

LANSOME II (W)

	J	MB	S	Z	Mean
		(±0.57)			(±0.28)
O	42.7	34.4	43.2	43.2	40.9
F	43.6	35.6	43.7	44.3	41.8
		(1) and (2)			(±0.33)
N1	33.3	27.1	35.9	35.5	33.0
N2	45.2	37.6	45.7	46.6	43.8
N3	50.9	40.2	48.7	49.0	47.2
Mean (±0.40)	43.1	35.0	43.4	43.7	41.3

Mean D.M. %: 84.2

- (1) (±0.67) For use in horizontal and diagonal comparisons only
- (2) (±0.66) For use in vertical and interaction comparisons only