

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/R&W/B/3 - Deep-drilled Urea & 'nitro-chalk' - Barley

Rothamsted Research

Rothamsted Research (1970) *69/R&W/B/3 - Deep-drilled Urea & 'nitro-chalk' - Barley*; Numerical Results Of The Field Experiments 1969, pp 263 - 264 - DOI:

<https://doi.org/10.23637/ERADOC-1-96>

BARLEY

(69/R/B/3 and 69/W/B/3)

Deeply drilled urea and 'Nitro-Chalk' Rothamsted (R) Pastures and Woburn (W) Stackyard A1, 1969.

Design: 4 blocks of 8 plots (randomisation restricted) plus 1 nil plot per block.

Area of each plot: 0.0161. Area harvested: 0.0107.

Treatments: No nitrogen (NO) and all combinations of:

1. Form of nitrogen: 'Nitro-Chalk' (C), urea (U).
2. Method of application: Injected by 'Tume' drill 3-4 inches deep in rows 5.5 inches apart (I), broadcast (B).
3. Nitrogen: 0.5 (N1), 1.0 (N2) cwt N.

Basal applications: 265 lb (0:20:20) combine drilled.

Weedkillers:-

Pastures (R): Paraquat at 0.5 lb ion in 25 gals 2,4-D at 8 oz and dichlorprop at 32 oz in 20 gals.

Stackyard A1 (W): Ioxynil octanoate, bromoxynil octanoate and the iso-octyl ester of dichlorprop ('Oxytril P' at 1 pint in 25 gals).

Cultivations, etc.:

Pastures (R): Paraquat applied: 11 Oct, 1968. Ploughed: 13 Nov. Urea and 'Nitro-Chalk' injected by 'Tume' drill: 28 Mar, 1969. Seed drilled at 145 lb, urea and 'Nitro-Chalk' broadcast: 29 Mar. 2,4-D/dichlorprop applied: 13 May. Combine harvested: 18 Aug. Variety: Zephyr. Previous crops: Potatoes 1967, barley 1968.

Stackyard A1 (W): Ploughed: 14 Aug, 1968. Ploughed second time: 15 Nov. Treatments applied, seed drilled at 145 lb: 3 Apr, 1969. Weedkiller applied: 16 May. Combine harvested: 8 Aug. Variety: Zephyr. Previous crops: Fallow 1967, 1968.

- NOTES: (1) The 'Tume' drill was used on all plots to prepare the seedbed, it being drawn idle through the B and NO plots.
(2) Percentage of N in grain was determined.

Standard errors per plot: Grain, cwt:

Pastures (R): 2.04 or 4.3% (21 d.f.)

Stackyard A1 (W): 4.53 or 13.7% (21 d.f.)

SUMMARY OF RESULTS

GRAIN, CWT

	I	B	N1	N2	Mean
PASTURES (R)					
	(± 0.72)				(± 0.51)
C	48.0	46.3	47.5	46.8	47.1
U	47.9	47.4	48.8	46.6	47.7
	(± 0.72)				(± 0.51)
		I	48.6	47.3	48.0
		B	47.6	46.1	46.8
Mean (± 0.51)			48.1	46.7	47.4

NO 36.4
 General mean: 46.2

STACKYARD A1 (W)

	(± 1.60)				(± 1.13)
C	34.0	32.4	30.6	35.9	33.2
U	32.7	32.8	31.9	33.6	32.7
	(± 1.60)				(± 1.13)
		I	32.4	34.3	33.4
		B	30.1	35.1	32.6
Mean (± 1.13)			31.3	34.7	33.0

NO 10.0
 General mean: 30.9

Mean D.M. %: Pastures (R): 79.6
 Stackyard A1 (W): 88.3