

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



---

### 69/R&W/BE/3 - *Rhizobium* Strains - Beans

#### Rothamsted Research

Rothamsted Research (1970) 69/R&W/BE/3 - *Rhizobium* Strains - Beans ; Numerical Results Of The Field Experiments 1969, pp 282 - 283 - DOI: <https://doi.org/10.23637/ERADOC-1-96>

### SPRING BEANS

(69/R/BE/3 and 69/W/BE/3)

The effect on yield of inoculating beans with different strains of *Rhizobium leguminosarum* - Rothamsted (R) Stackyard and Woburn (W) Horsepool 1969.

Design: 4 randomised blocks of 6 plots.

Area of each plot:

Stackyard (R): 0.0202. Area harvested: 0.0127.  
Horsepool (W): 0.0193. Area harvested: 0.0121.

Treatments: Seed inoculum (*Rhizobium leguminosarum*):-

None	(0)
Strain 1015	(1)
Strain 1028	(2)
Strain 1034	(3)
Strain 1038	(4)
Strain DP	(5)

Basal applications:

Stackyard (R): 360 lb (0:14:28) injected by 'Tune' drill.  
Horsepool (W): 450 lb (0:14:28) placement drilled.  
Weedkiller (both fields): Simazine at 1 lb in 20 gals (25 gals on Horsepool (W)).  
Insecticide (both fields): Demeton-s-methyl at 3.5 oz in 37 gals (30 gals on Horsepool (W)).

Cultivations, etc.:

Stackyard (R): Ploughed: 25 Oct, 1968. Basal PK compound injected: 10 Mar, 1969. Seed drilled at 200 lb, weedkiller applied: 27 Mar. Insecticide applied: 25 June. Combine harvested: 10 Sept. Variety: Tarvin. Previous crops: Barley 1967, fallow 1968.  
Horsepool (W): Ploughed: 2 Dec, 1968. Seed drilled at 200 lb: 1 Apr, 1969. Weedkiller applied: 4 Apr. Insecticide applied: 19 June. Combine harvested: 8 Sept. Variety: Maris Bead. Previous crops: Potatoes 1967, winter wheat 1968.

NOTE: Counts of *Rhizobium leguminosarum* in soil were made in March before sowing and in June from the rhizosphere. Nodules on roots were also counted.

Standard errors per plot. Grain, cwt:

Stackyard (R): 1.04 or 3.5% (15 d.f.)  
Horsepool (W): 1.13 or 5.4% (15 d.f.)

SUMMARY OF RESULTS

GRAIN:CWT

0	1	2	3	4	5	Mean
STACKYARD (R)						
(±0.52)						
30.2	29.9	30.0	30.1	29.6	29.5	29.9

Mean D.M.%: 83.8

HORSEPOOL (W)

(±0.56)

21.7	20.0	21.6	20.3	20.9	21.8	21.1
------	------	------	------	------	------	------

Mean D.M.%: 81.2