

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



### 69/R/RA/3 - Insecticides & Beneficial Insects - Spring Oilseed Rape

#### Rothamsted Research

Rothamsted Research (1970) *69/R/RA/3 - Insecticides & Beneficial Insects - Spring Oilseed Rape ; Numerical Results Of The Field Experiments 1969*, pp 293 - 294 - DOI:

<https://doi.org/10.23637/ERADOC-1-96>

SPRING OILSEED RAPE

(69/R/RA/3)

Insecticides and beneficial insects, West Barnfield I, 1969.

Design: 3 randomised blocks of 3 plots.

Area of each plot: 0.3214. Area harvested: 0.0459.

Treatments: None (O), azinphos-methyl applied as 27% wettable powder at 6 oz a.i. (A), Endosulfan applied as 35% wettable powder at 7.3 oz a.i. (E) in 60 gals.

Basal applications: 8 cwt (20:10:10) in seedbed. Weedkiller: Paraquat at 0.75 lb ion in 20 gals.

Cultivations, etc.: Paraquat applied: 1 Oct, 1968. Ploughed: 19 Nov. Basal NPK applied: 14 Apr, 1969. Seed drilled at 8 lb: 16 Apr. Insecticides applied: 27 June. Combine harvested: 28 Aug. Variety: Nilla. Previous crops: Winter wheat 1967, barley, 1968.

NOTE: Insect counts were made with a stationary suction sampler and a portable suction sampler. Visual counts were made of insects on flower heads and estimates made of the number of blind stalks.

Standard errors per plot.

Grain, cwt: 0.57 or 2.8% (4 d.f.)  
Yields of Fixed oil: 26.6 or 3.2% (4 d.f.)

SUMMARY OF RESULTS

O	A	E	Mean
GRAIN: CWT			
	(±0.33)		
19.9	20.6	21.0	20.5
% FIXED OIL			
38.7	38.5	38.5	38.6
YIELD OF FIXED OIL: LB			
	(±15.4)		
821	844	861	842

Mean D.M.%: 77.8