

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/R/G/1 - Anhydrous & Aqueous Ammonia - Grass

Rothamsted Research

Rothamsted Research (1970) *69/R/G/1 - Anhydrous & Aqueous Ammonia - Grass* ; Numerical Results Of The Field Experiments 1969, pp 321 - 322 - DOI: <https://doi.org/10.23637/ERADOC-1-96>

GRASS

(69/R/G/1)

Anhydrous and aqueous ammonia, Appletree 1969.

Design: 3 randomised blocks of 26 plots.

Area of each plot: 0.0092. Area harvested: 0.0054.

Treatments: None (NO) (2 plots per block) and all combinations of:-

1. Nitrogen fertiliser and time of application:

Applied in autumn:

Injected anhydrous ammonia (14 Nov)

IAA

Injected aqueous ammonia (13 Nov)

IQA

Applied in spring:

Injected anhydrous ammonia (28 Mar)

IAS

Injected aqueous ammonia (24 Mar)

IQS

Broadcast 'Nitro-Chalk':

Applied in 3 equal dressings

BD

Applied as single dressing

BS

2. N: 1 (N1), 2 (N2), 3 (N3), 4 (N4) cwt (total for the season).

Standard applications: 8.25 cwt (0:14:28) in winter. Weedkiller:

Applied to block I and spot-sprayed on docks and nettles in blocks II and III - 2,4-D ('Dicotox' at 6 pints in 40 gals).

Cultivations, etc.: Basal PK compound applied: 15 Nov, 1968. 'Nitro-Chalk' applied (treatment BS and first dressing of BD): 24 Mar, 1969.

Weedkiller applied: 28 Apr. Cut 3 times: 4 June, 6 Aug, 15 Oct.

'Nitro-Chalk' applied after first 2 cuts for BD treatment.

NOTES: (1) Grass samples were taken to determine dry matter and percentage of N, P and K. Percentage of Mg was determined in some samples.

(2) Growth of grass in summer showed that the anhydrous ammonia injector did not function properly in autumn and yields from this treatment have been omitted.

Standard errors per plot. Grass, dry matter, cwt.

1st cut: 2.40 or 6.2% (38 d.f.)

2nd cut: 2.36 or 9.1% (38 d.f.)

3rd cut: 0.82 or 14.4% (38 d.f.)

Total of 3 cuts: 3.34 or 4.7% (38 d.f.)

SUMMARY OF RESULTS

DRY MATTER: CWT

| | IQA | IAS | IQS | BD | BS | Mean |
|--------------|------|------|------|------|------|------|
| 1ST CUT | | | | | | |
| (±1.38) | | | | | | |
| (±0.62) | | | | | | |
| N1 | 41.2 | 37.0 | 40.6 | 33.9 | 39.5 | 38.5 |
| N2 | 38.6 | 36.9 | 39.0 | 39.4 | 39.0 | 38.6 |
| N3 | 42.6 | 38.2 | 40.5 | 39.4 | 37.6 | 39.7 |
| N4 | 40.9 | 37.1 | 40.2 | 40.9 | 37.0 | 39.2 |
| Mean (±0.69) | 40.8 | 37.3 | 40.1 | 38.4 | 38.3 | 39.0 |

NO: 19.8 (±0.98)

General mean (Excl IAA): 37.3

Mean D.M. %: 15.0
(All plots)

2ND CUT

| | | | | | | |
|--------------|------|------|------|------|------|------|
| (±1.36) | | | | | | |
| (±0.61) | | | | | | |
| N1 | 14.9 | 18.5 | 17.9 | 23.2 | 16.0 | 18.1 |
| N2 | 24.8 | 21.2 | 27.5 | 29.3 | 28.8 | 26.3 |
| N3 | 28.7 | 23.1 | 31.4 | 31.9 | 31.5 | 29.3 |
| N4 | 27.5 | 30.5 | 30.2 | 27.1 | 32.3 | 29.5 |
| Mean (±0.68) | 24.0 | 23.3 | 26.7 | 27.9 | 27.1 | 25.8 |

NO: 10.9 (±0.96)

General mean (Excl IAA): 24.4

Mean D.M. %: 21.8
(All plots)