

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/R&W/M/2 - Inocula & N - Beans and Spring Wheat - Mixed Crop

Rothamsted Research

Rothamsted Research (1970) *69/R&W/M/2 - Inocula & N - Beans and Spring Wheat - Mixed Crop* ;
Numerical Results Of The Field Experiments 1969, pp 327 - 332 - DOI:

<https://doi.org/10.23637/ERADOC-1-96>

INOCULA AND N

(69/R/M/2 and 69/W/M/2)

Beans and spring wheat - Rothamsted (R) Stackyard and Woburn (W)
Horsepool.

Design: 4 blocks of 4 plots split into 4 sub plots with confounding.

Area of each sub plot: 0.0011. Area harvested Stackyard (R): 0.0003.
Horsepool (W): 0.0003.

Treatments: All combinations of:-

- Whole plots: 1. Beans, seed inoculum (Rhizobium leguminosarum): None (O), ineffective strain 1019 (I), effective strain 1038 (E). Spring wheat (SW).
- Sub plots: 2. Fertiliser: Unmanured (U), 360 lb of compound fertiliser (0:14:28) plus 1 ton ground chalk Stackyard (R) 1.5 tons Horsepool (W), (M).
3. Nitrogen: None (O), 1.0 (N1), 2.0 (N2), 3.0 (N3) cwt N as 'Nitro-Chalk', half in the seedbed, and half top dressed.

Standard applications: Weedkiller: Simazine to beans at 1 lb in 40 gals.
Insecticide: Dimethoate to beans at 0.4 fl oz in 3 gals.

Cultivations, etc.:

Stackyard (R): Ploughed: 25 Oct, 1968. Ground chalk, PK and first half N applied: 27 Mar, 1969. Beans drilled at 200 lb, spring wheat drilled at 180 lb: 2 Apr. Weedkiller applied: 11 Apr. Remaining half N applied: 5 May. First harvest: 7 July. Insecticide applied: 30 July. Second harvest: 4 Aug. Varieties: Spring beans - Tarvin, spring wheat - Kolibri. Previous crops: Barley 1967, fallow 1968.

Horsepool (W): Ploughed: 4 Dec, 1968. Ground chalk, PK, and first half N applied: 8 Apr, 1969. Beans drilled at 200 lb, spring wheat drilled at 180 lb: 9 Apr. Weedkiller applied: 11 Apr. Remaining half N applied: 8 May. First harvest: 8 July. Insecticide applied: 30 July. Second harvest: 7 Aug. Varieties: Spring beans - Tarvin, spring wheat - Kolibri. Previous crops: Potatoes 1967, winter wheat 1968.

- NOTES: (1) Soil samples were taken for pH determination in February, and Rhizobium counts in February and June.
- (2) Grain yields were not taken - separate part plots were harvested green on two occasions.
- (3) Dry matter and % N determinations were made on crop produce.

Standard errors per plot: Dry matter, cwt:

Stackyard (R): 1st cut:	Whole plot:	1.22	or	3.2%	(6 d.f.)
	Sub plot:	5.20	or	13.4%	(24 d.f.)
2nd cut:	Whole plot:	5.33	or	9.3%	(6 d.f.)
	Sub plot:	6.76	or	11.8%	(24 d.f.)
Horsepool (W): 1st cut:	Whole plot:	3.66	or	7.6%	(6 d.f.)
	Sub plot:	5.88	or	12.3%	(24 d.f.)
2nd cut:	Whole plot:	5.04	or	7.5%	(6 d.f.)

SUMMARY OF RESULTS

STACKYARD (R)

DRY MATTER: CWT

	O	I	E	SW	Mean
	1ST CUT				
	(1) and (2)				(±0.92)
U	27.4	24.9	27.0	70.3	37.4
M	29.2	28.1	24.0	78.8	40.0
	(3) and (4)				(±1.30)
N0	29.4	23.7	24.5	47.4	31.2
N1	28.1	27.9	25.9	80.5	40.6
N2	27.0	26.5	26.1	84.8	41.1
N3	28.8	28.0	25.3	85.6	41.9
Mean (±0.61)	28.3	26.5	25.5	74.6	38.7

(1) (±1.44) (3) (±2.33) For use in horizontal and diagonal comparisons only

(2) (±1.84) (4) (±2.60) For use in vertical and interaction comparisons only

Mean D.M. %: 18.7

STACKYARD (R)

DRY MATTER: CWT

	O	I	E	SW	Mean
2ND CUT					
	(1) and (2)				(±1.19)
U	47.7	42.2	40.5	101.1	57.9
M	39.7	43.9	40.8	102.7	56.8
	(3) and (4)				(±1.69)
NO	41.8	35.6	39.6	74.5	47.9
N1	46.3	47.0	41.6	111.8	61.6
N2	42.9	48.7	39.5	105.6	59.2
N3	43.9	40.9	42.0	115.7	60.6
Mean (±2.67)	43.7	43.1	40.7	101.9	57.3

(1) (±3.16) (3) (±3.96) For use in horizontal and diagonal comparisons only
 (2) (±2.39) (4) (±3.38) For use in vertical and interaction comparisons only

Mean D.M. %: 26.8

HORSEPOOL (W)

DRY MATTER: CWT

	O	I	E	SW	Mean
	1ST CUT				
	(1) and (2)				(±1.04)
U	38.3	38.4	37.6	74.2	47.1
M	39.7	39.6	37.2	78.7	48.8
	(3) and (4)				(±1.47)
NO	35.8	36.9	35.0	59.4	41.8
N1	43.4	41.0	36.4	80.5	50.3
N2	37.0	37.1	38.2	82.2	48.6
N3	39.8	41.0	40.1	84.0	51.2
Mean (±1.83)	39.0	39.0	37.4	76.5	48.0

(1) (±2.35) (3) (±3.14) For use in horizontal and diagonal comparisons only
 (2) (±2.08) (4) (±2.94) For use in vertical and interaction comparisons only

Mean D.M. %: 18.5

HORSEPOOL (W)
DRY MATTER: CWT

	O	I	E	SW	Mean
2ND CUT					
(1) and (2)					
U	60.8	60.2	58.1	87.2	66.6
M	60.5	57.9	56.0	97.4	67.9
(3) and (4)					
NO	56.3	58.8	52.3	65.2	58.2
N1	65.0	61.1	56.4	97.4	70.0
N2	59.2	60.9	59.7	93.5	68.3
N3	62.3	55.3	59.6	113.1	72.6
Mean (± 2.52)	60.7	59.0	57.0	92.3	67.3

(1) (± 3.11) (3) (± 4.05) For use in horizontal and diagonal comparisons only
 (2) (± 2.58) (4) (± 3.65) For use in vertical and interaction comparisons only

Mean D.M. %: 30.8