

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1969

[Full Table of Content](#)



69/R/RN/7 - Residual Phosphate Rotation - Potatoes, Barley & Swedes

Rothamsted Research

Rothamsted Research (1970) *69/R/RN/7 - Residual Phosphate Rotation - Potatoes, Barley & Swedes ; Numerical Results Of The Field Experiments 1969*, pp 86 - 88 - DOI:

<https://doi.org/10.23637/ERADOC-1-96>

RESIDUAL PHOSPHATE ROTATION

(69/R/RN/7)

The long term and residual effects of phosphate fertilisers - Great Field IV and Sawyers I, the 10th year. For treatments and previous years' results see 'Details' 1967 and 'Results' 67/B/6 and 68/B/5.

Area of each plot:

Great Field IV: 0.0193.

Sawyers I: 0.0212.

Area harvested: Potatoes and barley: 0.0129. Swedes: 0.0096.

Area harvested: Potatoes and barley: 0.0141. Swedes: 0.0106.

Standard applications: Manures as previously. Weedkillers: Potatoes: Paraquat at 0.375 lb ion plus linuron at 0.75 lb ion in 37 gals. Fungicide: Mancozeb at 1.2 lb in 37 gals on 3 occasions. Insecticide: Demeton-s-methyl at 3.5 oz applied with second spraying with mancozeb. Haulm destroyer: Undiluted BOV at 15 gals. Barley: Weedkiller: 2,4-D at 8 oz and dichlorprop at 32 oz in 20 gals.

Cultivations, etc. (both fields): Ploughed: 21 Nov, 1968. Potatoes: Fertilisers applied: 14 Apr, 1969. Plots rotary cultivated, potatoes planted: 16 Apr. Paraquat and linuron applied: 13 May. Grubbed: 19 June. Rotary ridged: 25 June. Mancozeb applied: 15 July, 4 Aug, 21 Aug. Insecticide applied: 4 Aug. BOV applied: 4 Sept. Lifted: 23 Sept. Barley: Ground chalk applied at 23 cwt: 12 Nov, 1968. Fertilisers applied: 11 Mar, 1969. Seed drilled at 140 lb: 25 Mar. Weedkiller applied: 20 May. Combine harvested: 22 Aug. Swedes: Fertilisers applied: 29 Apr, 1969. Seed drilled at 1.25 lb: 14 May. Singled: 25 June. Lifted: 21 Oct.

Standard errors per plot.

Sawyers I:

Potatoes, total tubers, tons:	0.425 or 3.6% (11 d.f.)
Barley, grain, cwt:	2.17 or 5.2% (11 d.f.)
Swedes, fresh weight, tons:	1.059 or 10.2% (11 d.f.)

SUMMARY OF RESULTS

POTATOES

Treatment	TOTAL TUBERS: TONS		% WARE	
	Great Field IV Mean	Sawyers I Mean	Great Field IV Mean	Sawyers I Mean
		(±0.301)		
O	12.84	9.28	97.2	95.3
A1	14.68	11.03	96.6	94.9
A2	15.10	11.96	97.0	95.7
A3	16.28	12.50	96.8	95.2
A4	15.75	12.74	94.9	94.3
T1	15.79	12.08	96.4	95.2
T2	16.25	12.60	95.0	94.8
R2	14.82	11.51	97.1	96.5
R3	15.59	12.49	97.1	95.1
R4	15.74	12.76	95.1	94.7
G1	14.24	10.38	97.8	95.5
S1	13.71	10.81	96.6	95.1
Mean	15.07	11.68	96.5	95.2

BARLEY

	GRAIN: CWT		STRAW: CWT	
			(±1.53)	
O	30.1	39.1	27.3	28.5
A1	37.2	42.2	31.9	35.0
A2	31.4	44.7	33.9	34.7
A3	33.8	42.9	30.8	38.2
A4	34.9	42.3	28.0	34.3
T1	32.8	43.3	35.3	35.8
T2	40.1	43.0	35.0	34.8
R2	36.5	41.9	36.4	36.3
R3	33.5	44.3	46.0	37.3
R4	35.9	42.7	34.3	37.4
G1	31.7	40.1	28.3	30.5
S1	31.8	39.0	29.0	27.9
Mean	34.1	42.1	33.0	34.2
Mean D.M. %:	80.4	83.8	88.6	91.6

SWEDES, ROOTS: TONS

Treat- ment	Great Field IV Mean	Sawyers I Mean
		(±0.749)
O	5.70	2.78
A1	10.93	8.97
A2	16.67	12.06
A3	17.64	13.72
A4	15.56	14.44
T1	14.72	11.98
T2	16.48	11.83
R2	15.46	10.99
R3	16.58	12.00
R4	16.07	13.26
G1	10.00	5.24
S1	10.37	7.26
Mean	13.85	10.38