

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/W/RN/15 Rotation and Fumigation - Barley, Potatoes, Sugar Beet

Rothamsted Research

Rothamsted Research (1971) *70/W/RN/15 Rotation and Fumigation - Barley, Potatoes, Sugar Beet ; Numerical Results Of The Field Experiments 1970*, pp 124 - 128 - DOI:
<https://doi.org/10.23637/ERADOC-1-59>

ROTATION AND FUMIGATION

(70/W/RN/15)

The effects of soil fumigants on a rotation of crops - Woburn Butt Close 1970, 2nd year.

For design, treatments and previous year's results see 'Results' 69/W/RN/15.

Area of each sub-plot: 0.0048. Area harvested: Barley, potatoes: 0.0013, sugar beet: 0.0038.

NOTE: One of the sub-plots of each plot, untreated in 1969, is now used to test dazomet at 200 lb before all crops (AZ).

Standard applications:

Barley: Manures: 280 lb (0:20:20). Weedkiller: Ioxynil at 7.5 oz plus mecoprop at 22.5 oz in 25 gals.

Potatoes: Manures: 940 lb (0:14:28). Weedkiller: Linuron at 1.0 lb plus paraquat at 0.37 lb ion in 33 gals. Fungicide: Mancozeb at 1.2 lb in 37 gals on two occasions. Insecticide: Demeton-s-methyl at 3.5 fl oz in 37 gals on one occasion with fungicide.

Sugar beet: Manures: 940 lb (0:14:28). Weedkiller: Phenmedipham ('Betanal' at 5 pts in 20 gals). Insecticide: Demeton-s-methyl at 3.5 fl oz in 25 gals.

Cultivations, etc.:

Barley: Ploughed: 12 Nov, 1969. Dazomet applied and rotary cultivated in: 14 Nov. DD injected: 24 Nov. Reploughed: 3 Mar, 1970. N applied, power harrowed, seed combine drilled at 140 lb: 27 Mar. Weedkiller applied: 16 May. Combine harvested: 26 Aug. Variety: Julia.

Potatoes: Ploughed: 12 Nov, 1969. Dazomet applied and rotary cultivated in: 14 Nov. DD injected: 24 Nov. Reploughed: 3 Mar, 1970. N applied: 3 Apr. PK applied: 14 Apr. Rotary cultivated, potatoes planted: 22 Apr. Weedkiller applied: 13 May. Grubbed: 3 June. Rotary ridged: 5 June. Fungicide plus insecticide applied: 31 July. Fungicide applied: 18 Aug. Sprayed with undiluted BOV at 15 gals: 19 Sept. Haulm destroyed mechanically: 23 Sept. Lifted: 30 Sept. Variety: Pentland Crown.

Sugar beet: Rotary cultivated twice: 11 Sept, 1969 and 30 Sept. Dazomet applied and rotary cultivated in: 14 Nov. DD injected: 24 Nov. Ploughed: 3 Mar, 1970. N applied: 3 Apr. PK applied: 14 Apr. Power harrowed, seed drilled at 5 lb: 20 Apr. Weedkiller applied: 15 May. Singled: 26 May. Insecticide applied: 22 June. Lifted: 26 Oct. Variety: Klein E.

70/W/RN/15

- NOTES: (1) Soil samples were taken from all N2 plots for eelworm counts, (*Heterodera rostochiensis*, *H.avenae*) in late November.
(2) The yields of barley grain and straw have been adjusted for a linear trend down the whole plots.

Standard errors per sub-plot.

Barley, grain, cwt:	3.33 or 25.8% (17 d.f.)
Sugar beet, roots tons:	1.252 or 9.1% (18 d.f.)
Total sugar, cwt:	4.60 or 9.4% (18 d.f.)
Potatoes, total tubers, tons:	2.012 or 11.2% (18 d.f.)

Barley straw, grain, sugar beet, potatoes, and eelworm counts are given in the tables on pages 125-128.

70/W/RN/15

SUMMARY OF RESULTS

BARLEY

	O,S&R	P	B	A	AZ	Mean
GRAIN: CWT						
	(±1.36)			(±2.36)*		
N1	7.2	12.2	8.9	10.1	16.9	9.9
N2	10.6	15.7	15.9	18.4	16.7	14.1
N3	12.5	18.0	15.7	14.0	17.7	14.7
Mean	10.1 (±0.78)	15.3	13.5 (±1.36)	14.2	17.1	12.9

* For use in horizontal and interaction comparisons only

STRAW: CWT

N1	8.7	10.5	11.2	11.1	14.0	10.4
N2	12.3	15.6	15.2	16.5	16.0	14.3
N3	11.4	14.2	14.5	12.6	16.5	13.1
Mean	10.8	13.4	13.6	13.4	15.5	12.6

Mean D.M. %: Grain: 80.1
Straw: 71.8

70/W/RN/15

POTATOES

	O,B&R	P	S	A	AZ	Mean
TOTAL TUBERS: TONS						
	(±0.821)			(±1.423)*		
N1	14.79	15.15	17.23	16.23	17.02	15.71
N2	18.07	19.79	19.36	20.75	19.71	19.12
N3	16.83	19.40	20.31	22.14	20.01	18.91
Mean	16.56 (±0.474)	18.12	18.97 (±0.822)	19.71	18.91	17.91

* For use in horizontal and interaction comparisons only

% WARE: 1.5 INCH

N1	98.2	98.8	98.4	98.1	98.5	98.4
N2	98.4	98.1	98.7	98.1	98.7	98.4
N3	98.0	98.2	98.3	98.4	98.3	98.2
Mean	98.2	98.4	98.5	98.2	98.5	98.3

70/W/RN/15

SUGAR BEET

	O, P&R	S	B	A	AZ	Mean
ROOTS (WASHED): TONS						
	(±0.511)			(±0.885)*		
N1	10.85	10.46	11.41	10.66	15.73	11.54
N2	14.34	14.92	13.58	16.07	15.52	14.73
N3	14.02	14.90	17.01	15.74	16.70	15.20
Mean	13.07 (±0.295)	13.43	14.00 (±0.511)	14.16	15.99	13.82
SUGAR %						
N1	18.4	17.9	18.0	18.0	18.2	18.2
N2	17.7	17.7	17.5	17.1	18.0	17.6
N3	17.3	16.9	17.3	16.5	17.1	17.1
Mean	17.8	17.5	17.6	17.2	17.8	17.7
TOTAL SUGAR: CWT						
	(±1.88)			(±3.25)*		
N1	40.0	37.5	40.9	38.3	57.6	42.0
N2	50.9	52.7	47.6	54.9	55.9	52.0
N3	48.6	50.4	58.9	51.9	57.3	52.0
Mean	46.5 (±1.08)	46.9	49.2 (±1.88)	48.4	56.9	48.7

* For use in horizontal and interaction comparisons only