

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/R/CS/6 Wheat After Intensive Barley -w. Wheat

Rothamsted Research

Rothamsted Research (1971) *70/R/CS/6 Wheat After Intensive Barley -w. Wheat* ; Numerical Results Of The Field Experiments 1970, pp 133 - 134 - DOI: <https://doi.org/10.23637/ERADOC-1-59>

WHEAT AFTER INTENSIVE BARLEY

(70/R/CS/6)

The tenth year of the 'Intensive Barley Growing Experiment', Little Knott I. For design, treatments etc. see 'Results' 61/C/8 and 69/R/CS/6.

Area of each plot: 0.0212. Area harvested: 0.0140.

Cropping: Winter wheat was grown over the whole experiment, with the exception of sequence 1 (winter wheat 1969, preceded by 6 years of continuous barley), which was fallowed.

Basal applications: 780 lb (0:14:28) ploughed in, 220 lb (0:14:28) combine drilled. Weedkillers: Paraquat at 0.75 lb ion in 20 gals. Terbutryne and related triazines ('Prebane' at 4 lb in 20 gals). 2,4-D at 0.5 lb and dichlorprop at 2 lb in 20 gals.

Cultivations, etc.: First basal PK applied: 16 Sept, 1969. Paraquat applied: 17 Sept. Ploughed: 23 Sept. All plots rotary cultivated: 10 Nov. Seed combine drilled at 160 lb: 11 Nov. 'Prebane' applied: 14 Nov. 2,4-D/dichlorprop applied: 5 May, 1970. 'Nitro-Chalk' applied: 13 May. Combine harvested: 26 Aug. Variety: Joss Cambier.

- NOTES: (1) Yields were taken for all sequences except sequence 1 (fallow).
(2) Estimates of take-all (*Ophiobolus graminis*) were made in early July.
(3) For previous years' results see 'Results' 61/C/8, 62/C/7, 63/C/2, 64/C/2, 65/C/2, 66/C/2, 67/C/2, 68/C/2, 69/R/CS/6.

Standard error per plot.

Grain, cwt: 3.94 or 11.5% (35 d.f.)

70/R/CS/6

SUMMARY OF RESULTS

GRAIN: CWT

Crop Sequences

Crop in 1961	2	3	4	5	6	7	8	9	10	
	WS	O	BE	WS	WS	B	WS	WS	BE	
2	O	WS	O	BE	WS	B	WS	WW	WW	
3	BE	O	WS	O	BE	B	WS	WW	P	
4	B	BE	O	WS	O	B	WS	WW	B	
5	B	B	BE	O	WS	B	WS	WW	BE	
6	B	B	B	BE	O	B	WS	WW	WW	
7	B	B	B	B	BE	B	WS	WW	P	
8	B	B	B	B	B	B	WS	F	B	
9	WW	WW	WW	WW	WW	WW	WW	WW	F	Mean
					(±2.79)					(±0.93)
N3	30.0	32.3	29.0	24.7	22.0	32.9	29.2	32.4	47.5	31.1
N5	28.5	30.8	33.0	32.5	25.7	34.2	32.0	34.9	49.4	33.4
N7	30.2	41.9	31.6	32.8	29.2	35.4	39.2	40.6	48.8	36.6
N9	36.1	33.4	26.9	33.8	31.7	39.2	34.8	38.9	53.8	36.5
Mean (±1.39)	31.2	34.6	30.1	30.9	27.2	35.4	33.8	36.7	49.9	34.4

Mean D.M. %: 83.4