

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



## 70/R/CS/23 Simulated Grazing - Old Grass

### Rothamsted Research

Rothamsted Research (1971) *70/R/CS/23 Simulated Grazing - Old Grass* ; Numerical Results Of The Field Experiments 1970, pp 162 - 165 - DOI: <https://doi.org/10.23637/ERADOC-1-59>

SIMULATED GRAZING

(70/R/CS/23)

Old grass, plot 6 Park Grass 1970 the fourth year. For treatments, etc., and for previous years' results see 'Results' 67/C/38, 68/C/29 and 69/R/CS/23.

Treatments G+H and HG+ were replaced by treatments G+ (cuttings returned after each cut) and G- (cuttings removed). The new treatments were allocated at random for each N level in each block.

Cultivations, etc.: Basal P, K, Na and Mg applied: 4 Dec, 1969.

The following treatments were cut on the dates shown:-

24 Apr, 1970:	G+	G-	G-H		
11 May:				HG-	H
12 May:	G+	G-	G-H		
1 June:	G+	G-		HG-	
18 June:	G+	G-		HG-	
30 June:	G+	G-	G-H	HG-	H
21 July:	G+	G-	G-H		
4 Aug:	G+	G-	G-H		
2 Sept:	G+	G-	G-H	HG-	H
16 Sept:	G+	G-		HG-	
5 Oct:	G+	G-		HG-	
26 Oct:	G+	G-	G-H	HG-	H

Calcium nitrate applied on 16 Mar, 12 May, 30 June, 2 Sept.

NOTES: (1) On the G+ plots some grass returned at previous cutting was recovered by the rotary mower. Therefore these yields were overestimated.

(2) The percentages of N, P and K in the dry grass were measured.

Standard errors per plot. Dry matter, cwt:

1st Period (cuts 1-2):	3.53 or 22.7% (28 d.f.)
2nd Period (cuts 3-5):	3.18 or 19.6% (28 d.f.)
3rd Period (cuts 6-8):	2.76 or 15.8% (28 d.f.)
4th Period (cuts 9-11):	1.26 or 14.8% (28 d.f.)
Total of all 4 Periods (cuts 1-11):	4.65 or 8.0% (28 d.f.)

70/R/GS/23

SUMMARY OF RESULTS

DRY MATTER: CWT

1ST PERIOD

Treatment	H	H(G-)	G-(H)	G+	G-	Mean
Cut with	M	M	R	R	R	
			(±2.04)			(±0.91)
NO	3.3	2.3	5.4	6.0	6.7	4.7
N1	20.7	18.7	13.8	15.7	16.8	17.1
N2	29.8	29.6	20.6	23.3	20.4	24.7
Mean (±1.18)	18.0	16.9	13.3	15.0	14.6	15.5

R plots cut twice.

2ND PERIOD

Treatment	H	(H)G-	(G-)H	G+	G-	Mean
Cut with	M	R	M	R	R	
			(±1.84)			(±0.82)
NO	14.4	9.6	16.6	17.7	10.2	13.7
N1	17.5	11.6	18.5	28.8	11.7	17.6
N2	15.7	10.3	22.3	25.1	12.8	17.2
Mean (±1.06)	15.9	10.5	19.1	23.9	11.6	16.2

R plots cut three times.

M = Motor scythe

R = Rotary mower

( ) = Not applied in this period

70/R/CS/23

DRY MATTER: CWT

3RD PERIOD

Treatment	H	H(G-)	G-(H)	G+	G-	Mean
Cut with	M	M	R	R	R	
			(±1.60)			(±0.71)
NO	11.9	12.3	11.8	7.9	7.4	10.3
N1	18.4	22.2	14.2	20.3	13.0	17.6
N2	26.4	31.9	20.6	22.7	21.5	24.6
Mean (±0.92)	18.9	22.1	15.5	17.0	14.0	17.5

R plots cut three times

4TH PERIOD

Treatment	H	(H)G-	(G-)H	G+	G-	Mean
Cut with	M	R	M	R	R	
			(±0.73)			(±0.32)
NO	4.9	6.4	5.4	6.6	4.9	5.7
N1	10.4	7.1	9.3	15.0	7.7	9.9
N2	9.3	6.7	12.4	13.4	7.7	9.9
Mean (±0.42)	8.2	6.7	9.0	11.7	6.8	8.5

R plots cut three times

70/R/CS/23

DRY MATTER: CWT

TOTAL OVER ALL PERIODS

Treatment Cut with	H	HG-	G-H	G+	G-	Mean	Under same management	
	M	M&R	M&R	R	R		H(G-)	G-(H)
			(±2.68)			(±1.20)		
NC	34.6	30.6	39.2	38.2	29.2	34.4	36.6	33.2
N1	67.0	59.6	55.8	79.8	49.3	62.3	68.7	46.7
N2	81.2	78.6	75.9	84.5	62.4	76.5	96.2	58.2
Mean (±1.55)	60.9	56.2	57.0	67.5	47.0	57.7	67.1	46.0