



P, K AND TAKE-ALL

(70/R/CS/24)

P, K and take-all (*Ophiobolus graminis*), West Barnfield II 1970, the third year, barley. For treatments etc. and for the previous years' results see 'Results' 68/C/16 and 69/R/CS/24.

Design: 2 replicates of 5x2x4 in 4 blocks of 10 plots split into 2 for N.

Area of each sub plot: 0.0127. Area harvested: 0.0067.

N test: A test of 0.3 (N1), 0.6 (N2), 0.9 (N3), 1.2 (N4) cwt N as 'Nitro-Chalk' is now applied to half plots. The test is cumulative in successive years.

Basal applications: Manures: None. Weedkiller: 2,4-D at 0.5 lb and dichlorprop at 2 lb in 20 gals.

Cultivations, etc.: Ploughed: 29 Sept, 1969. N, P and K applied: 25 Mar, 1970. Seed drilled at 140 lb: 31 Mar. Weedkiller applied: 18 May. Combine harvested: 15 Aug. Variety: Julia.

NOTE: Estimates of eyespot (*Cercospora herpotrichoides*) and take-all (*Ophiobolus graminis*) were made in early July.

Standard errors per plot.

Grain, cwt: Whole plot: 1.91 or 5.7% (17 d.f.)  
Sub plot: 2.21 or 6.6% (20 d.f.)  
Pooled (used for calculation of standard errors in summary): 2.45 or 7.3% (37 d.f.)

70/R/CS/24

SUMMARY OF RESULTS

GRAIN: CWT

|              | P0   | P1   | P4      | P6*  | P24* | Mean    |
|--------------|------|------|---------|------|------|---------|
|              |      |      | (±0.86) |      |      | (±0.39) |
| K1           | 24.4 | 33.0 | 38.0    | 31.7 | 37.7 | 33.0    |
| K4           | 24.5 | 31.3 | 38.6    | 34.2 | 40.3 | 33.8    |
|              |      |      | (±1.22) |      |      | (±0.55) |
| N1           | 24.4 | 30.8 | 35.4    | 32.0 | 36.6 | 31.8    |
| N2           | 23.0 | 32.3 | 40.2    | 31.9 | 39.7 | 33.4    |
| N3           | 26.3 | 33.6 | 39.6    | 34.0 | 40.2 | 34.7    |
| N4           | 24.2 | 32.2 | 37.9    | 33.9 | 39.5 | 33.5    |
| Mean (±0.61) | 24.5 | 32.2 | 38.3    | 33.0 | 39.0 | 33.4    |
|              | N1   | N2   | N3      | N4   |      |         |
|              |      |      | (±0.77) |      |      |         |
| K1           | 31.9 | 33.2 | 33.9    | 33.0 |      |         |
| K4           | 31.8 | 33.7 | 35.6    | 34.0 |      |         |

\* Applied 1968 only

Mean D.M. %: 86.1

70/R/GS/24

GRAIN: CWT

|    | K1      |      |      |      |      | K4   |      |      |      |      |
|----|---------|------|------|------|------|------|------|------|------|------|
|    | P0      | P1   | P4   | P6   | P24  | P0   | P1   | P4   | P6   | P24  |
|    | (±1.73) |      |      |      |      |      |      |      |      |      |
| N1 | 24.2    | 31.5 | 34.4 | 31.9 | 37.4 | 24.5 | 30.1 | 36.4 | 32.1 | 35.8 |
| N2 | 25.1    | 32.9 | 40.5 | 30.1 | 37.2 | 20.9 | 31.6 | 39.9 | 33.7 | 42.3 |
| N3 | 22.7    | 34.1 | 38.9 | 33.4 | 40.2 | 29.8 | 33.0 | 40.2 | 34.7 | 40.1 |
| N4 | 25.6    | 33.8 | 38.1 | 31.5 | 36.1 | 22.7 | 30.6 | 37.7 | 36.3 | 42.8 |

70/R/CS/24

STRAW: CWT

|      | P0   | P1   | P4   | P6*  | P24* | Mean |
|------|------|------|------|------|------|------|
| K1   | 9.2  | 11.7 | 14.8 | 11.8 | 13.9 | 12.3 |
| K4   | 10.8 | 13.5 | 17.9 | 14.4 | 17.3 | 14.8 |
| N1   | 9.4  | 11.4 | 12.7 | 12.0 | 13.7 | 11.8 |
| N2   | 8.9  | 13.1 | 17.8 | 13.1 | 15.7 | 13.7 |
| N3   | 11.5 | 14.1 | 16.7 | 13.1 | 15.9 | 14.3 |
| N4   | 10.1 | 11.9 | 18.2 | 14.4 | 16.9 | 14.3 |
| Mean | 10.0 | 12.6 | 16.3 | 13.1 | 15.6 | 13.5 |
|      | N1   | N2   | N3   | N4   |      |      |
| K1   | 11.3 | 12.4 | 13.1 | 12.3 |      |      |
| K4   | 12.3 | 15.1 | 15.4 | 16.3 |      |      |

\* Applied 1968 only

Mean D.M. %: 88.8

70/R/CS/24

STRAW: CWT

|    | K1  |      |      |      |      | K4   |      |      |      |      |
|----|-----|------|------|------|------|------|------|------|------|------|
|    | PO  | P1   | P4   | P6   | P24  | PO   | P1   | P4   | P6   | P24  |
| N1 | 9.8 | 11.3 | 11.2 | 11.2 | 13.0 | 8.9  | 11.4 | 14.2 | 12.7 | 14.4 |
| N2 | 9.1 | 12.1 | 15.9 | 10.8 | 14.0 | 8.7  | 14.2 | 19.7 | 15.3 | 17.4 |
| N3 | 8.9 | 12.7 | 16.2 | 12.4 | 15.1 | 14.2 | 15.4 | 17.2 | 13.7 | 16.7 |
| N4 | 8.9 | 10.7 | 15.7 | 12.8 | 13.3 | 11.3 | 13.1 | 20.6 | 16.0 | 20.6 |