

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Numerical Results of the Field Experiments 1970

[Full Table of Content](#)



70/W/CS/28 Fumigants and Irrigation - Barley

Rothamsted Research

Rothamsted Research (1971) *70/W/CS/28 Fumigants and Irrigation - Barley* ; Numerical Results Of The Field Experiments 1970, pp 173 - 175 - DOI: <https://doi.org/10.23637/ERADOC-1-59>

FUMIGANTS AND IRRIGATION

(70/W/CS/28)

The residual effects of fumigants for the control of *Pratylenchus* spp. in barley - Woburn Butt Close Series III, 3rd year. For design, treatments etc. and previous years' results see 'Results' 68/C/34 and 69/W/CS/28.

NOTE: Fumigant treatments are residual from 1968. Irrigation and nitrogen treatments are cumulative 1968, 1969 and 1970.

Area of each sub plot: 0.0050. Area harvested: 0.0020.

Basal applications: 280 lb (0:20:20). Weedkiller: Ioxynil at 7.5 oz and mecoprop at 22.5 oz in 25 gals.

Cultivations, etc.: Ploughed: 27 Aug, 1969. Power harrowed: 1 Nov. Deep-tine cultivated: 3 Nov. Power harrowed: 26 Mar, 1970. Seed combine drilled at 140 lb: 27 Mar. 'Nitro-Chalk' applied: 3 Apr. Weedkiller applied: 16 May. Irrigation applied at 1.0 inch on five occasions: 9 June, 11 June, 16 June, 22 June, 7 July. Combine harvested: 26 Aug. Variety: Maris Badger.

NOTE: Crop and soil samples were taken at intervals through the growing season for nematode counts.

Standard errors per plot. Grain, cwt:
Whole plot: 1.81 or 7.1% (6 d.f.)
Quarter plot: 2.26 or 8.9% (24 d.f.)
Eighth plot: 2.80 or 11.0% (32 d.f.)

70/W/CS/28

SUMMARY OF RESULTS

GRAIN: CWT

1968

	O	C	D	T	Mean
1968 - 70	(1) and (2)				(±0.74)
O	15.6	21.9	23.5	17.6	19.7
I	30.3	30.4	32.9	30.8	31.1
1968 - 70					
	NO	N1	N2	N3	
	(3) and (4)				
O	9.2	22.1	24.3	23.2	
I	12.9	34.1	42.3	35.3	
1968	(5) and (6)				(±0.65)
O	10.4	24.5	31.6	25.4	23.0
C	12.0	26.9	35.7	30.1	26.2
D	11.0	32.7	36.1	32.9	28.2
T	10.8	28.0	29.7	28.5	24.2
Mean (±0.57)	11.0	28.1	33.3	29.2	25.4

Mean D.M. %: 85.0

- (1) (±1.09) (3) (±1.02) (5) (±1.19) For use in vertical and diagonal comparisons only
 (2) (±0.92) (4) (±0.81) (6) (±1.14) For use in horizontal and interaction comparisons only

70/W/CS/28

STRAW: CWT

1968

	O	C	D	T	Mean
1968 - 70					
O	10.9	13.9	15.2	12.0	13.0
I	24.9	24.0	27.1	24.7	25.2
1968 - 70					
	NO	N1	N2	N3	
O	6.1	14.0	16.1	15.9	
I	8.8	24.5	34.6	32.8	
1968					
O	7.4	18.0	25.6	20.7	17.9
C	7.8	16.8	26.6	24.5	18.9
D	7.7	23.2	27.0	26.7	21.1
T	6.9	18.9	22.2	25.3	18.3
Mean	7.4	19.2	25.4	24.3	19.1

Mean D.M. %: 87.5